



Research Note

RN/13/11

Parametric Completeness for Separation Theories

June 7, 2013

James Brotherston Jules Villard

Abstract

In this paper, we address the logical gap between provability in the logic BBI, which is the standard propositional basis for *separation logic*, and validity in an intended class of *separation models*, as employed in applications of separation logic such as program verification. An intended class of separation models is usually specified by a collection of axioms describing various useful model properties, which we call a *separation theory*.

Our three main contributions are as follows. First, we show that several important properties of separation models are not definable in BBI. Second, we show that these properties become definable when a simple theory of naming is added to BBI, in the manner familiar from *hybrid logic*. We initially present a simple, binder-free hybrid version of BBI and then later add a binder, which can capture more complex model properties. Third, we present an axiomatic proof system for our hybrid logic whose extension with any set of syntactically “pure” axioms is sound and complete with respect to the models satisfying those axioms. As a corollary of this general result, we obtain sound and complete axiomatic proof systems for a large class of separation theories which, to the best of our knowledge, includes all those appearing in the published literature.

Our results are obtained by adaptations of techniques from the general setting of modal and hybrid logics.

Parametric Completeness for Separation Theories

James Brotherston* Jules Villard†

Dept. of Computer Science, University College London, UK

Abstract

In this paper, we address the logical gap between provability in the logic BBI, which is the standard propositional basis for *separation logic*, and validity in an intended class of *separation models*, as employed in applications of separation logic such as program verification. An intended class of separation models is usually specified by a collection of axioms describing various useful model properties, which we call a *separation theory*.

Our three main contributions are as follows. First, we show that that several important properties of separation models are not definable in BBI. Second, we show that these properties become definable when a simple theory of naming is added to BBI, in the manner familiar from *hybrid logic*. We initially present a simple, binder-free hybrid version of BBI and then later add a binder, which can capture more complex model properties. Third, we present an axiomatic proof system for our hybrid logic whose extension with any set of syntactically “pure” axioms is sound and complete with respect to the models satisfying those axioms. As a corollary of this general result, we obtain sound and complete axiomatic proof systems for a large class of separation theories which, to the best of our knowledge, includes all those appearing in the published literature.

Our results are obtained by adaptations of techniques from the general setting of modal and hybrid logics.

Categories and Subject Descriptors F.3.1 [Logics and Meanings of Programs]: Specifying and Verifying and Reasoning about Programs—Logics of programs

General Terms Theory, verification

Keywords Bunched logic, separation logic, hybrid logic

1. Introduction

Essentially, all models are wrong, but some are useful.

G.E.P. Box and N.R. Draper [4], 1987

In mathematical logic, there is a notable tension between *provability* in a logic — which captures *validity* in some general class of models of that logic — and *validity in the intended model(s)* of practical or theoretical interest. For example, the axioms of Peano

* Research supported by an EPSRC Career Acceleration Fellowship.

† Research supported by EPSRC grant EP/H008373/2.

arithmetic (PA) hold in many nonstandard models as well as the usual standard model of the natural numbers \mathbb{N} [18]. Thus, as is well known, there are many true statements of arithmetic over \mathbb{N} that do not hold in every model of the Peano axioms, and so are not provable in PA. This *incompleteness* of logical proof systems with respect to a particular model of interest is sometimes an unavoidable phenomenon. For example, as is well known, true arithmetic is not recursively enumerable, so cannot be captured by any finitary proof system. In other cases, it might happen that completeness for intended models can be gained by adding (perhaps infinitely many) axioms to the proof system, at the expense of simplicity and perhaps of desirable structural properties such as cut-elimination. Thus, when formulating axiomatisations of a theory, some trade-off is usually inevitable between logical generality on the one hand, and completeness for the intended model(s) on the other.

In this paper, we consider this trade-off in the context of *separation logic*, an established formalism for reasoning about heap-manipulating programs [7, 24, 26]. The purely propositional part of separation logic is usually considered to be given by Boolean BI (from now on BBI), which is a particular flavour of *bunched logic* obtained by freely combining the connectives of multiplicative intuitionistic linear logic with those of standard classical logic [17, 23]. Provability in BBI corresponds to validity in the general class of relational commutative monoids [14]. Applications of separation logic, on the other hand, typically deal with *specific* such (classes of) models, based on the composition of heaps (see [6] for a survey of the models used in practice). Unsurprisingly, these heap models exhibit various interesting mathematical properties that are not true of all relational commutative monoids, and thus are not captured by provability in BBI. For example, composition of heaps is a cancellative partial (binary) function, which is a special case of the ternary relation in a relational commutative monoid. Various collections of such properties have previously been advanced in the literature as a suitable abstraction over concrete heap models [9, 13]. We list these properties in Definition 3.1, and call a given collection of such properties a *separation theory*. Our aim is to develop logics in which provability corresponds more closely to validity in the class of models determined by a separation theory.

In this paper, we make three main contributions:

- First, we show in section 3 that BBI is insufficiently expressive to accurately capture most separation theories. Specifically, we show that several commonly considered model properties are not definable by any BBI-formula (partial functionality and cancellativity of the composition being two of the most prominent examples). Undefinability of a property means that the logic is fundamentally incapable of distinguishing models with the property from those without it.

- Second, we introduce in section 4 a simple *hybrid* extension HyBBI of BBI, which bears the same relation to BBI as normal hybrid logic does to normal modal logic (see [1–3] for an

overview). That is, HyBBI extends BBI with a theory of *naming*: we introduce a second sort of atoms, called *nominals*, which are interpreted as individual states in a model; and we also add a unary hybrid modality $@_\ell$ (parameterised by the nominal ℓ), so that the hybrid formula $@_\ell A$ is satisfied at any world in a model just when A is satisfied at the world denoted by the nominal ℓ .

Despite the simplicity of this extension, which is conservative over standard BBI, the hybrid logic HyBBI is expressive enough to define most of the separation theories we consider. However, for more complex model properties such as the *cross-split* property of [13], still more expressivity is required: we show in Section 7 how to gain this expressivity by adding the \downarrow binder of hybrid logic to HyBBI.

- Third, we provide a Hilbert-style axiomatic proof system for HyBBI that is *parametrically* sound and complete with respect to any given separation theory. That is, whenever the proof system is extended with the axioms defining a separation theory, the resulting extension is sound and complete with respect to the class of models determined by that theory. Although such axiomatic proof systems are not very suitable for practical reasoning, they provide a very useful proof-theoretic characterisation of validity which can be used as a baseline, e.g., for tableau or sequent-style proof systems.

We present the axiom system and its soundness result in Section 5, and give the completeness theorem in Section 6. The extension of these results to the case of HyBBI with the \downarrow binder is given in Section 7.

All of our technical results are obtained by adaptations of techniques from modal and hybrid logic. This should come as no surprise, since bunched logics can quite straightforwardly be seen as modal logics. Indeed, this view has been exploited previously in the literature, e.g., to obtain completeness results for various bunched logics [5, 8]. However, as far as we know, HyBBI represents the first explicit application of hybrid logic to the setting of bunched / separation logic. In particular, previous work on hybrid logics has seemingly been confined to modal logic with unary modalities connected by de Morgan duality, whereas in this setting we consider the case of binary modalities connected by residuation.

Interestingly, the key concept from hybrid logic, i.e. the explicit naming of elements in the underlying model, has been used implicitly several times in the literature on the proof theory of BBI. For example, the labelled tableau system for BBI provided in [20], which was recently proven complete for partial functional BBI-models [19], relies on a system of semantic labels which pick out individual model states in much the same way as nominal atoms in hybrid logic. Even more recently, labelled nested [22] and non-nested [16] sequent calculi for BBI have appeared, employing semantic labels in a broadly similar way. While such works add names or labels to proof systems as auxiliary tools for simplifying proof search in standard BBI, here we consider these features to be first class components of the logic. Indeed, we believe that it should be reasonably straightforward to adapt the labelled proof systems in the literature to yield cut-free proof theories for our hybrid extensions of BBI.

2. Syntax and semantics of BBI

In this section, we give the usual definitions of formulas of BBI and their Kripke semantics, given by relational commutative monoids (see e.g. [14]).

Definition 2.1 (BBI-formula). Let \mathcal{V} be a denumerably infinite set of *propositional variables*. BBI-formulas are built from propositional variables $P \in \mathcal{V}$ using the usual connectives ($\top, \perp, \neg, \wedge, \vee, \rightarrow$)

of classical logic, and the so-called “multiplicative” connectives, consisting of the constant I and binary operators $*$ and \multimap .

We assume that \neg has the highest precedence, followed by $*$, \wedge and \vee , with \rightarrow and \multimap having lowest precedence.

Definition 2.2 (BBI frames and models). A BBI-frame is a tuple $\langle W, \circ, E \rangle$, where $\circ : W \times W \rightarrow \mathcal{P}(W)$ and $E \subseteq W$. We extend \circ pointwise to $\mathcal{P}(W) \times \mathcal{P}(W) \rightarrow \mathcal{P}(W)$ by

$$W_1 \circ W_2 \stackrel{\text{def}}{=} \bigcup_{w_1 \in W_1, w_2 \in W_2} w_1 \circ w_2$$

A BBI-frame $\langle W, \circ, E \rangle$ is a BBI-model if \circ is commutative and associative, and $w \circ E = \{w\}$ for all $w \in W$.

Definition 2.3 (BBI-validity). Let $M = \langle W, \circ, E \rangle$ be a BBI-frame. A *valuation* for M is a function ρ that assigns to each propositional variable $P \in \mathcal{V}$ a set $\rho(P) \subseteq W$. Given any valuation ρ for M , any $w \in W$ and any BBI-formula A , we define the forcing relation $M, w \models_\rho A$ by induction on A :

$$\begin{aligned} M, w \models_\rho P &\Leftrightarrow w \in \rho(P) \\ M, w \models_\rho \top &\Leftrightarrow \text{always} \\ M, w \models_\rho \perp &\Leftrightarrow \text{never} \\ M, w \models_\rho \neg A &\Leftrightarrow M, w \not\models_\rho A \\ M, w \models_\rho A_1 \wedge A_2 &\Leftrightarrow M, w \models_\rho A_1 \text{ and } M, w \models_\rho A_2 \\ M, w \models_\rho A_1 \vee A_2 &\Leftrightarrow M, w \models_\rho A_1 \text{ or } M, w \models_\rho A_2 \\ M, w \models_\rho A_1 \rightarrow A_2 &\Leftrightarrow M, w \models_\rho A_1 \text{ implies } M, w \models_\rho A_2 \\ M, w \models_\rho I &\Leftrightarrow w \in E \\ M, w \models_\rho A_1 * A_2 &\Leftrightarrow \exists w_1, w_2 \in W. w \in w_1 \circ w_2 \text{ and } \\ &\quad M, w_1 \models_\rho A_1 \text{ and } M, w_2 \models_\rho A_2 \\ M, w \models_\rho A_1 \multimap A_2 &\Leftrightarrow \forall w', w'' \in W. \text{ if } w'' \in w \circ w' \text{ and } \\ &\quad M, w' \models_\rho A_1 \text{ then } M, w'' \models_\rho A_2 \end{aligned}$$

A is said to be *valid* in M if $M, w \models_\rho A$ for any valuation ρ and for all $w \in W$. A is *valid* if it is valid in all BBI-models.

Definition 2.4. We define \mathbf{K}_{BBI} to be the proof system obtained by extending a complete Hilbert system for classical logic with the following axioms and inference rules for $*$, \multimap and I (where $A \vdash B$ is syntactic sugar for the formula $A \rightarrow B$):

$$\begin{array}{c} A * B \vdash B * A \quad A * (B * C) \vdash (A * B) * C \\ \\ A \vdash A * I \quad A * I \vdash A \\ \\ \frac{A_1 \vdash B_1 \quad A_2 \vdash B_2}{A_1 * A_2 \vdash B_1 * B_2} \quad \frac{A * B \vdash C}{A \vdash B \multimap C} \quad \frac{A \vdash B \multimap C}{A * B \vdash C} \end{array}$$

The following result is due to Galmiche and Larchey-Wendling [14].

Theorem 2.5. A BBI-formula is \mathbf{K}_{BBI} -provable if and only if it is BBI-valid.

3. Definable and undefinable properties in BBI

In this section, we review a number of interesting properties of BBI-models encountered in the literature on separation logic, and consider the problem of whether or not these properties can be expressed, or *defined*, by formulas of BBI. Specifically, we show that some important such properties are *not* definable in BBI, essentially because they are not generally preserved over morphisms between BBI-models that preserve validity.

Definition 3.1 (Separation theories). Letting $M = \langle W, \circ, E \rangle$ be a BBI-model, we introduce the following properties of interest:

- Partial functionality*: $w, w' \in w_1 \circ w_2$ implies $w = w'$;
- Cancellativity*: $(w \circ w_1) \cap (w \circ w_2) \neq \emptyset$ implies $w_1 = w_2$;
- Single unit*: $|E| = 1$, i.e. $w, w' \in E$ implies $w = w'$;
- Indivisible units*: $(w \circ w') \cap E \neq \emptyset$ implies $w \in E$;

Disjointness: $w \circ w \neq \emptyset$ implies $w \in E$;

*Divisibility:*¹ for every $w \notin E$ there are $w_1, w_2 \notin E$ such that $w \in w_1 \circ w_2$;

Cross-split property: whenever $(t \circ u) \cap (v \circ w) \neq \emptyset$, there exist tv, tw, uv, uw such that $t \in tv \circ tw, u \in uv \circ uw, v \in tv \circ uv$ and $w \in tw \circ uw$.

Any given collection of model properties from the above list is called a *separation theory*.

Various different separation theories have been considered in the literature on separation logic. For example, a BBI-model that is both partial functional and cancellative is called a *separation algebra* in [13], while in [9] the same term defines a BBI-model that is partial functional and cancellative with a single unit, and, in the views framework [12], the same term again refers to a BBI-model that is simply partial functional. The significance of the other properties is explained in detail in [13].

In the following, we abuse notation slightly by identifying a property of BBI-models with the class of BBI-models satisfying that property.

Definition 3.2 (Definability). Given a language \mathcal{L} of formulas, a property \mathcal{P} of BBI-models is said to be \mathcal{L} -*definable* if there exists an \mathcal{L} -formula A such that for all BBI-models M ,

$$A \text{ is valid in } M \iff M \in \mathcal{P}.$$

We remark that definability could as well be defined on BBI-frames, not just BBI-models. However, our main concern is with properties of BBI-models (and in particular those listed in Definition 3.1). Note that the property of being a BBI-model, among all frames, is itself BBI-definable: take as the defining formula the conjunction of the top four axioms in Definition 2.4 (which define associativity, commutativity and the unit law $E \circ w = \{w\}$).

Proposition 3.3. *The indivisible units property and the divisibility property are both BBI-definable, as follows:*

$$\begin{array}{ll} \text{Indivisible units:} & I \wedge (A * B) \vdash A \\ \text{Divisibility:} & \neg I \vdash \neg I * \neg I \end{array}$$

Proof. The case of the indivisible units property is shown in [6]. For the case of divisibility, we proceed as follows:

(\Leftarrow) Assume that M is divisible, let ρ be a valuation for M and let $w \in W$. To show that (sp) is valid, we suppose that $M, w \models_{\rho} \neg I$, i.e., that $w \notin E$, and require to show that $M, w \models_{\rho} (\neg I) * (\neg I)$. Divisibility gives us $w_1, w_2 \in W \setminus E$ such that $w \in w_1 \circ w_2$; thus, $M, w \models_{\rho} (\neg I) * (\neg I)$.

(\Rightarrow) Assume that (sp) is valid in M , and suppose that $w \in W \setminus E$. Then, $M, w \models_{\rho} \neg I$, hence by validity of (sp) we have $M, w \models_{\rho} (\neg I) * (\neg I)$. This gives us w_1, w_2 such that $w \in w_1 \circ w_2$ where $M, w_1 \models_{\rho} \neg I$ and $M, w_2 \models_{\rho} \neg I$, i.e. $w_1, w_2 \notin E$ as required. \square

We now develop some tools, based on the usual ones from modal logic [1], for showing that a given property is *not* definable in BBI. Our main such tool is a *bounded morphism*, which is a structure-preserving function between BBI-frames.

Definition 3.4 (Bounded morphic image). Let $M = \langle W, \circ, E \rangle$ and $M' = \langle W', \circ', E' \rangle$ be BBI-models. A *bounded morphism* from M to M' is a function $f : W \rightarrow W'$ satisfying the following:

1. $w \in E$ iff $f(w) \in E'$;
2. $w \in w_1 \circ w_2$ implies $f(w) \in f(w_1) \circ' f(w_2)$;

¹Called ‘‘splittability’’ in [13].

3. $f(w) \in w'_1 \circ' w'_2$ implies $\exists w_1, w_2 \in W. w \in w_1 \circ w_2$ and $f(w_1) = w'_1$ and $f(w_2) = w'_2$;
4. $w'_2 \in f(w) \circ' w'_1$ implies $\exists w_1, w_2 \in W. w_2 \in w \circ w_1$ and $f(w_1) = w'_1$ and $f(w_2) = w'_2$;

We say M' is a *bounded morphic image* of M , written $M \rightarrow M'$, if there is a surjective bounded morphism from M to M' .

Lemma 3.5. *Let M and M' be BBI-models with $M \rightarrow M'$. Then any BBI-formula valid in M is also valid in M' .*

Proof. We write $M = \langle W, \circ, E \rangle$ and $M' = \langle W', \circ', E' \rangle$, and let $f : W \rightarrow W'$ be a surjective bounded morphism from M to M' . Suppose for contradiction that A is valid in M , but not in M' . Thus there exists a valuation ρ' for M' and $w' \in W'$ such that $M', w' \not\models_{\rho'} A$. We define a valuation ρ for M as follows:

$$\rho(P) \stackrel{\text{def}}{=} \{w \in W \mid f(w) \in \rho'(P)\}$$

As f is surjective, there is a $w \in W$ such that $w' = f(w)$. To obtain the required contradiction, we claim that $M, w \not\models_{\rho} A$. To show this claim, we prove by structural induction on A that for all $w \in W$, we have $M, w \models_{\rho} A$ if and only if $M', f(w) \models_{\rho'} A$. We just show one of the cases for the classical connectives, as the others are either similar, or else trivial.

Case $A = P \in \mathcal{V}$. Using the definition of ρ , we have as required:

$$\begin{aligned} M, w \models_{\rho} A &\iff w \in \rho(P) \\ &\iff f(w) \in \rho'(P) \\ &\iff M', f(w) \models_{\rho'} P \end{aligned}$$

Case $A = B \rightarrow C$. We proceed as follows:

$$\begin{aligned} M, w \models_{\rho} B \rightarrow C &\iff M, w \models_{\rho} B \text{ implies } M, w \models_{\rho} C \\ &\iff M', f(w) \models_{\rho'} B \text{ implies } M', f(w) \models_{\rho'} C \quad (\text{by ind. hyp.}) \\ &\iff M', f(w) \models_{\rho'} B \rightarrow C \end{aligned}$$

Case $A = I$. Using condition 1 in Defn. 3.4, we have as required:

$$M, w \models_{\rho} I \iff w \in E \iff f(w) \in E' \iff M', f(w) \models_{\rho'} I$$

*Case $A = B * C$.* We show each implication separately.

(\Rightarrow) Supposing that $M, w \models_{\rho} B * C$, we have $w \in w_1 \circ w_2$ with $M, w_1 \models_{\rho} B$ and $M, w_2 \models_{\rho} C$. By induction hypothesis, $M', f(w_1) \models_{\rho'} B$ and $M', f(w_2) \models_{\rho'} C$. Furthermore, using condition 2 in Defn. 3.4, we have $f(w) \in f(w_1) \circ' f(w_2)$. Thus $M', f(w) \models_{\rho'} B * C$ as required.

(\Leftarrow) Supposing that $M', f(w) \models_{\rho'} B * C$, we have $f(w) \in w'_1 \circ' w'_2$ with $M', w'_1 \models_{\rho'} B$ and $M', w'_2 \models_{\rho'} C$. By condition 3 in Defn. 3.4, there are $w_1, w_2 \in W$ with $w \in w_1 \circ w_2$ and $f(w_1) = w'_1$ and $f(w_2) = w'_2$. Thus, by the induction hypothesis, we have $M, w_1 \models_{\rho} B$ and $M, w_2 \models_{\rho} C$. Hence $M, w \models_{\rho} B * C$ as required.

Case $A = B \multimap C$. We show each implication separately.

(\Rightarrow) Suppose $M, w \models_{\rho} B \multimap C$. To show that $M', f(w) \models_{\rho'} B \multimap C$, we assume that $w'_2 \in f(w) \circ' w'_1$ and $M', w'_1 \models_{\rho'} B$, and must show $M', w'_2 \models_{\rho'} C$. By condition 4 in Defn. 3.4, there are $w_1, w_2 \in W$ with $w_2 \in w \circ w_1$ and $f(w_1) = w'_1$ and $f(w_2) = w'_2$. Thus, by the induction hypothesis, $M, w_1 \models_{\rho} B$. Since $M, w \models_{\rho} B \multimap C$, we obtain $M, w_2 \models_{\rho} C$, which yields the required $M', w'_2 \models_{\rho'} C$ using the induction hypothesis.

(\Leftarrow) Suppose $M', f(w) \models_{\rho'} B \multimap C$. To show that $M, w \models_{\rho} B \multimap C$, we assume that $w_2 \in w \circ w_1$ and $M, w_1 \models_{\rho} B$, and

must show $M, w_2 \models_{\rho} C$. By condition 2 in Defn. 3.4, we have $f(w_2) \in f(w) \circ f(w_1)$, and by induction hypothesis we have $M', f(w_1) \models_{\rho'} B$. Since $M', f(w) \models_{\rho'} B \multimap C$, we obtain $M', f(w_2) \models_{\rho'} C$, which then yields the required $M, w_2 \models_{\rho} C$ using the induction hypothesis. This completes all cases. \square

Lemma 3.6. *Let \mathcal{P} be a property of BBI-models, and let M, M' be BBI-models such that $M \in \mathcal{P}$, $M' \notin \mathcal{P}$ and $M \rightarrow M'$. Then \mathcal{P} is not BBI-definable.*

Proof. Suppose for contradiction that there exists a BBI-formula A valid in exactly those BBI-models with property \mathcal{P} . Then, in particular, A is valid in M . Since $M \rightarrow M'$, we have by Lemma 3.5 that A must be valid in M' , and hence $M' \in \mathcal{P}$, contradicting the assumption that $M' \notin \mathcal{P}$. \square

In particular, Lemma 3.6 implies that if $M \rightarrow M'$ and M is a BBI-model, then so is M' (otherwise the class of all BBI-models would not be BBI-definable among all BBI-frames, contradiction).

Theorem 3.7. *Partial functionality is not BBI-definable.*

Proof. By Lemma 3.6, it suffices to exhibit a pair of BBI-models M and M' such that M is partial functional, M' is not partial functional and $M \rightarrow M'$. We define BBI-models $M = \langle W, \circ, E \rangle$ and $M' = \langle W', \circ', E' \rangle$ as follows:

$$\begin{aligned} W &\stackrel{\text{def}}{=} \{e, v_1, v_2, x_1, x_2, y, z\} & E &\stackrel{\text{def}}{=} \{e\} \\ w \circ e &= e \circ w \stackrel{\text{def}}{=} \{w\} \text{ for all } w \in W \\ x_1 \circ v_1 &= v_1 \circ x_1 \stackrel{\text{def}}{=} \{y\} & x_1 \circ v_2 &= v_2 \circ x_1 \stackrel{\text{def}}{=} \{y\} \\ x_2 \circ v_1 &= v_1 \circ x_2 \stackrel{\text{def}}{=} \{z\} & x_2 \circ v_2 &= v_2 \circ x_2 \stackrel{\text{def}}{=} \{z\} \\ W' &\stackrel{\text{def}}{=} \{e, v, x, y, z\} & E' &\stackrel{\text{def}}{=} \{e\} \\ w \circ' e &= e \circ' w \stackrel{\text{def}}{=} \{w\} \text{ for all } w \in W' \\ x \circ' v &= v \circ' x \stackrel{\text{def}}{=} \{y, z\} \end{aligned}$$

with $w_1 \circ w_2 = w_1 \circ' w_2 \stackrel{\text{def}}{=} \emptyset$ for all other w_1 and w_2 .

First, we verify that M and M' are indeed BBI-models. Commutativity and the unit law hold in both models by construction. Associativity of \circ and \circ' is straightforward to check since $w_1 \circ (w_2 \circ w_3)$ and $w_1 \circ' (w_2 \circ' w_3)$ are always empty unless one of w_1, w_2, w_3 is e .

Next, we note that M is partial functional since $|w_1 \circ w_2| \leq 1$ for all $w_1, w_2 \in W$ by construction, whereas M' is not partial functional since $z, y \in x \circ' v$ but $z \neq y$.

Finally, we claim that $M \rightarrow M'$, i.e., that there is a surjective bounded morphism from M to M' . Define $f : W \rightarrow W'$ by:

$$\begin{aligned} f(v_1) &= f(v_2) \stackrel{\text{def}}{=} v & f(x_1) &= f(x_2) \stackrel{\text{def}}{=} x \\ f(w) &\stackrel{\text{def}}{=} w & (w \in \{e, u, v, y\}) \end{aligned}$$

Clearly f is surjective, so it just remains to check the 4 bounded morphism conditions in Definition 3.4:

1. Trivial, since $E = E' = \{e\}$ and $f(e) = e$.
2. We just check that every membership statement in the definition of \circ maps under f to a corresponding membership statement in the definition of \circ' . E.g., since $y \in x_1 \circ v_2$, we need to check that $f(y) \in f(x_1) \circ' f(v_2)$, i.e., $y \in x \circ' v$, which is the case.
3. We need to check that every membership statement $f(w) \in w'_1 \circ w'_2$ in the definition of \circ' can be “traced back” under f to a corresponding membership statement in the definition of \circ . E.g., since $f(z) \in x \circ' v$, we need w_1, w_2 such that $z \in w_1 \circ w_2$ and $f(w_1) = x, f(w_2) = v$. By taking, say, $w_1 = x_2$ and $w_2 = v_2$, we are done.

4. Similar to item 3 above, but for membership statements of the form $w'_2 \in f(w) \circ w'_1$. E.g., since $y \in f(v_2) \circ' x$, we need w_1, w_2 such that $w_2 \in v_2 \circ w_1$ and $f(w_1) = x, f(w_2) = y$. By taking $w_1 = x_1, w_2 = y$ we are done. This completes the proof. \square

At this juncture it is perhaps worth noting that there is no *a priori* connection between definability of a property on one hand, and the existence of complete proof systems for models having the property on the other. In particular, Theorem 3.7 says nothing about the existence of proof theories for BBI that are complete for partial functional models. In fact, Larchey-Wendling and Galmiche exhibit in [21] a formula that shows that \mathbf{K}_{BBI} is incomplete for such models. What Theorem 3.7 shows in addition is that, if one were to add enough (perhaps infinitely many) axioms to \mathbf{K}_{BBI} so as to obtain a complete system for partial functional models, then provability in this system still would not exclude all strictly relational models. (One can contrast this situation, e.g., with that of \mathbf{K}_{BBI} 's commutativity axiom $A * B \vdash B * A$, which is easily seen to define commutativity and therefore to exclude all non-commutative models.)

Theorem 3.8. *Cancellativity is not BBI-definable.*

Proof. By Lemma 3.6, it suffices to exhibit a pair of BBI-models M and M' such that M is cancellative, M' is not cancellative and $M \rightarrow M'$. We define BBI-models $M = \langle W, \circ, E \rangle$ and $M' = \langle W', \circ', E' \rangle$ as follows:

$$\begin{aligned} W &\stackrel{\text{def}}{=} \{e, v_1, v_2, x, y, z_1, z_2\} & E &\stackrel{\text{def}}{=} \{e\} \\ w \circ e &= e \circ w \stackrel{\text{def}}{=} \{w\} \text{ for all } w \in W \\ v_1 \circ x &= x \circ v_1 \stackrel{\text{def}}{=} \{z_1\} & v_2 \circ x &= x \circ v_2 \stackrel{\text{def}}{=} \{z_2\} \\ v_1 \circ y &= y \circ v_1 \stackrel{\text{def}}{=} \{z_2\} & v_2 \circ y &= y \circ v_2 \stackrel{\text{def}}{=} \{z_1\} \\ W' &\stackrel{\text{def}}{=} \{e, x, v, y, z\} & E' &\stackrel{\text{def}}{=} \{e\} \\ w \circ' e &= e \circ' w \stackrel{\text{def}}{=} \{w\} \text{ for all } w \in W' \\ v \circ' x &= x \circ' v = v \circ' y = y \circ' v \stackrel{\text{def}}{=} \{z\} \end{aligned}$$

First, it is straightforward to verify that M and M' are indeed BBI-models, with associativity holding because $w_1 \circ (w_2 \circ w_3) \neq \emptyset$ implies one of w_1, w_2, w_3 is e (and similarly for \circ').

Next, we note that M is cancellative because, by construction, $w' \in (w \circ w_1) \cap (w \circ w_2)$ implies $w_1 = w_2 = e$. On the other hand, M' is not cancellative, for $z \in (v \circ' x) \cap (v \circ' y)$ but $x \neq y$.

Finally, we need a surjective bounded morphism from M to M' . We define a map $f : W \rightarrow W'$ by

$$\begin{aligned} f(v_1) &= f(v_2) \stackrel{\text{def}}{=} v & f(z_1) &= f(z_2) \stackrel{\text{def}}{=} z \\ f(w) &\stackrel{\text{def}}{=} w & (w \in \{e, x, y\}) \end{aligned}$$

The verification that f is indeed a surjective bounded morphism is similar to that in the proof of Theorem 3.7. \square

We note that Theorem 3.8 in fact maps a model that is both partial functional and cancellative with a single unit to a non-cancellative model. Thus, it also establishes that neither the class of models that are partial functional *and* cancellative (the “separation algebras” of [13]) nor the subclass of such models having a single unit (the “separation algebras” of [9]) are BBI-definable.

Theorem 3.9. *Disjointness is not BBI-definable.*

Proof. By Lemma 3.6, it suffices to exhibit a pair of BBI-models M and M' such that M is disjoint, M' is not and $M \rightarrow M'$. We define BBI-models $M = \langle W, \circ, E \rangle$ and $M' = \langle W', \circ', E' \rangle$ as

follows:

$$\begin{aligned} W &\stackrel{\text{def}}{=} \{e, x, y\} & E &\stackrel{\text{def}}{=} \{e\} \\ w \circ e &= e \circ w \stackrel{\text{def}}{=} \{w\} \text{ for all } w \in W \\ x \circ y &= y \circ x \stackrel{\text{def}}{=} \{x, y\} \end{aligned}$$

$$\begin{aligned} W' &\stackrel{\text{def}}{=} \{e, x\} & E' &\stackrel{\text{def}}{=} \{e\} \\ w \circ' e &= e \circ' w \stackrel{\text{def}}{=} \{w\} \text{ for all } w \in W \\ x \circ' x &\stackrel{\text{def}}{=} \{x\} \end{aligned}$$

with $w_1 \circ w_2 = w_1 \circ' w_2 \stackrel{\text{def}}{=} \emptyset$ for all other w_1 and w_2 .

Similar to the previous Theorems 3.7 and 3.8, we can easily verify that M and M' are indeed BBI-models, with M disjoint by construction, whereas M' is not disjoint since $x \neq e$ and $x \circ' x \neq \emptyset$. We define a surjective bounded morphism f from M to M' by

$$f(e) \stackrel{\text{def}}{=} e \quad f(x) \stackrel{\text{def}}{=} x \quad f(y) \stackrel{\text{def}}{=} x$$

It just remains to check the bounded morphism conditions, which is similar to the verifications in Theorems 3.7 and 3.8. \square

Definition 3.10 (Disjoint union). Let $M_1 = \langle W_1, \circ_1, E_1 \rangle$, $M_2 = \langle W_2, \circ_2, E_2 \rangle$ be BBI-models, where W_1 and W_2 are disjoint sets. Then $M_1 \uplus M_2$, the *disjoint union* of M_1 and M_2 is defined as

$$M_1 \uplus M_2 \stackrel{\text{def}}{=} \langle W_1 \cup W_2, \circ_1 \cup \circ_2, E_1 \cup E_2 \rangle$$

Lemma 3.11. *Let M_1, M_2 be BBI-models. Then any BBI-formula valid in both M_1 and M_2 is also valid in $M_1 \uplus M_2$.*

Proof. We write $M_1 = \langle W_1, \circ_1, E_1 \rangle$ and $M_2 = \langle W_2, \circ_2, E_2 \rangle$. Suppose for contradiction that A is valid in M_1 and M_2 , but not in $M_1 \uplus M_2$. Thus there exists a valuation ρ and $w \in W_1 \cup W_2$ such that $M_1 \uplus M_2, w \not\models_\rho A$. Without loss of generality, we assume $w \in W_1$ (the other case is similar). We define a valuation for M_1 by

$$\rho_1(P) \stackrel{\text{def}}{=} \rho(P) \cap M_1$$

To obtain the required contradiction, we claim that $M_1, w \not\models_{\rho_1} A$ (contradicting the supposition that A is valid in M_1). To show this claim, we prove by structural induction on A that for all $w \in W_1$, we have $M_1, w \models_{\rho_1} A$ if and only if $M_1 \uplus M_2, w \models_\rho A$. We omit the cases for the classical connectives, since they are trivial or immediate by induction hypothesis.

Case $A = P$. Using the fact that $w \in W_1$ by assumption, we have $w \in \rho_1(P)$ iff $w \in \rho(P)$, i.e. $M_1, w \models_{\rho_1} P$ iff $M_1 \uplus M_2, w \models_\rho P$ as required.

Case $A = I$. Since E_1 and E_2 are disjoint, we have $w \in E_1$ iff $w \in E_1 \cup E_2$, i.e. $M_1, w \models_{\rho_1} I$ iff $M_1 \uplus M_2, w \models_\rho I$, as required.

*Case $A = B * C$.* We proceed as follows:

$$\begin{aligned} &M_1 \uplus M_2, w \models_\rho B * C \\ \Leftrightarrow &\exists w_1, w_2 \in W_1 \cup W_2. w \in w_1(\circ_1 \cup \circ_2)w_2 \text{ and} \\ &M_1 \uplus M_2, w_1 \models_\rho B \text{ and } M_1 \uplus M_2, w_2 \models_\rho C \\ \Leftrightarrow &\exists w_1, w_2 \in W_1. w \in w_1 \circ_1 w_2 \text{ and } M_1 \uplus M_2, w_1 \models_\rho B \\ &\text{and } M_1 \uplus M_2, w_2 \models_\rho C \quad (\text{since } W_1, W_2 \text{ disjoint}) \\ \Leftrightarrow &\exists w_1, w_2 \in W_1. w \in w_1 \circ_1 w_2 \text{ and } M_1, w_1 \models_{\rho_1} B \text{ and} \\ &M_1, w_2 \models_{\rho_1} C \quad (\text{by ind. hyp.}) \\ \Leftrightarrow &M_1, w \models_{\rho_1} B * C \end{aligned}$$

Case $A = B \multimap C$. Similar to the case $A = B * C$. \square

Lemma 3.12. *Let \mathcal{P} be a property of BBI-models, and suppose that there exist BBI-models M_1 and M_2 such that $M_1, M_2 \in \mathcal{P}$ but $M_1 \uplus M_2 \notin \mathcal{P}$. Then \mathcal{P} is not BBI-definable.*

Proof. Suppose for contradiction that there exists a BBI-formula valid in exactly those BBI-models $M \in \mathcal{P}$. Thus, by assumption, A is valid in M_1 and in M_2 . By Lemma 3.11, A is valid in $M_1 \uplus M_2$, whence $M_1 \uplus M_2 \in \mathcal{P}$, contradicting the premise that $M_1 \uplus M_2 \notin \mathcal{P}$. \square

Note that, similar to Lemma 3.6, Lemma 3.12 implies that if M_1 and M_2 are BBI-models then so must be $M_1 \uplus M_2$ (otherwise BBI-models would again be undefinable among all BBI-frames).

Theorem 3.13. *The single-unit property is not BBI-definable.*

Proof. Let $M_1 = \langle \mathbb{N}, +, \{0\} \rangle$ and $M_2 = \langle \mathbb{N}', +', \{0'\} \rangle$ be disjoint, isomorphic copies of the monoid of natural numbers under addition. M_1 and M_2 are both single-unit BBI-models. However, $M_1 \uplus M_2$ is not a single-unit model, as its set of units is $\{0, 0'\}$. Thus, by Lemma 3.12, the single-unit property cannot be BBI-definable. \square

There is one property from Definition 3.1 that we have not yet considered: the cross-split property. In Section 7, we show that this property is definable in a relatively expressive hybrid extension of BBI including binders (see Proposition 7.3). The complication of expressing the property even in that logic leads us to strongly suspect it is not definable in BBI.

Conjecture 3.14. *The cross-split property is not BBI-definable.*

Cross-split is seemingly preserved by bounded morphic images, disjoint unions and by *generated submodels* (cf. [1]). We believe it should be possible to show its undefinability in BBI by employing a model construction based on *ultrafilter extensions* (cf. [1]), but that is beyond the scope of the present paper.

4. HyBBI: a basic hybrid extension of BBI

In this section, we present a hybrid extension of BBI, called HyBBI, based upon a simple fragment of *hybrid logic* [1–3]. Essentially, compared to standard BBI, our extension allows us to refer to individual elements of the underlying BBI-model (as opposed to sets of elements, as denoted by BBI-formulas). We demonstrate that this simple extension of the logic enables us to define the separation theory properties shown in the previous section to be undefinable in BBI.

Definition 4.1 (HyBBI-formula). We assume a fixed, denumerably infinite set \mathcal{N} of *nominals*, disjoint from the propositional variables. (We write lower case letters j, k, ℓ etc. for nominals to distinguish them from propositional variables.) A HyBBI-formula is defined as a BBI-formula (Defn. 2.1), except that (a) any nominal $\ell \in \mathcal{N}$ counts as an atomic HyBBI-formula, and (b) if A is a HyBBI-formula and ℓ a nominal then $@_\ell A$ is a HyBBI-formula.

A HyBBI-formula is said to be *pure* if it contains no propositional (or formula) variables.

Definition 4.2 (HyBBI-validity). A *hybrid valuation* ρ for a BBI-model $M = \langle W, \circ, E \rangle$ extends a standard valuation (see Defn. 2.3) by additionally mapping every nominal $\ell \in \mathcal{N}$ to an element $\rho(\ell) \in W$. Given any hybrid valuation ρ for M , any $w \in W$ and a HyBBI-formula A , we define the forcing relation $M, w \models_\rho A$ by extending the definition of the forcing relation in Defn. 2.3 by the following clauses for nominals and the $@_\ell$ modality:

$$\begin{aligned} M, w \models_\rho \ell &\Leftrightarrow w = \rho(\ell) \\ M, w \models_\rho @_\ell A &\Leftrightarrow M, \rho(\ell) \models_\rho A \end{aligned}$$

A is then said to be *valid in M* if $M, w \models_\rho A$ for all hybrid valuations ρ and all $w \in W$ (and simply *valid* if it is valid in all BBI-models).

We observe that HyBBI is a *conservative extension* of BBI; that is, every BBI-formula A that is valid according to Definition 2.3 is also valid according to Definition 4.2 (because the forcing relations in the two definition coincide on BBI-formulas). Thus every property of BBI-models definable in BBI is also definable in HyBBI. However, HyBBI is strictly more expressive than BBI: several properties not definable in BBI become definable in HyBBI.

Theorem 4.3. *The following properties from Definition 3.1 are HyBBI-definable via pure formulas:*

<i>Functionality:</i>	$\@_{\ell}(j * k) \wedge \@_{\ell'}(j * k) \vdash \@_{\ell} \ell'$	(pfn)
<i>Cancellativity:</i>	$\ell * j \wedge \ell * k \vdash \@_j k$	(cnc)
<i>Single unit:</i>	$\@_{\ell_1} I \wedge \@_{\ell_2} I \vdash \@_{\ell_1} \ell_2$	(su)
<i>Indivisible units:</i>	$I \wedge (\ell_1 * \ell_2) \vdash \ell_1$	(iu)
<i>Disjointness:</i>	$\ell * \ell \vdash I \wedge \ell$	(dis)
<i>Divisibility:</i>	$\neg I \vdash \neg I * \neg I$	(div)

Proof. The case of divisibility is already covered by Proposition 3.3, and the case of indivisible units is a straightforward modification of the argument in [6] (using nominal variables rather than propositional variables). We treat the other properties individually.

Functionality. (\Leftarrow) Assume M is partial functional, let ρ be a valuation for M and let $w \in W$. To show that (pfn) is valid in M , we assume that $M, w \Vdash_{\rho} \@_{\ell}(j * k) \wedge \@_{\ell'}(j * k)$, and must show that $M, w \Vdash_{\rho} \@_{\ell} \ell'$, i.e., that $\rho(\ell) = \rho(\ell')$.

By assumption, we have $\rho(\ell) \in \rho(j) \circ \rho(k)$ and $\rho(\ell') \in \rho(j) \circ \rho(k)$, hence by partial functionality of M we have $\rho(\ell) = \rho(\ell')$ as required.

(\Rightarrow) Assume that (pfn) is valid in M , and suppose $w', w \in w_1 \circ w_2$. We require to show that $w = w'$.

Since (pfn) is valid in M , we have $M, w \Vdash_{\rho} (\text{pfn})$ for all $w \in W$ and for any hybrid valuation ρ . We define a hybrid valuation ρ for M as follows:

$$\rho(\ell) = w \quad \rho(\ell') = w' \quad \rho(j) = w_1 \quad \rho(k) = w_2$$

Then, showing that $w = w'$ means showing that $M, \rho(\ell) \Vdash_{\rho} \ell'$, i.e., that $M, w \Vdash_{\rho} \@_{\ell} \ell'$. As (pfn) is valid in M , it suffices to show that $M, w \Vdash_{\rho} \@_{\ell}(j * k) \wedge \@_{\ell'}(j * k)$. Since $w, w' \in w_1 \circ w_2$ by assumption and $M, w_1 \Vdash_{\rho} j$ and $M, w_2 \Vdash_{\rho} k$ by construction, we obtain that $M, w \Vdash_{\rho} j * k$ and that $M, w' \Vdash_{\rho} j * k$, from which the result follows.

Cancellativity. (\Leftarrow) Assume M is cancellative, let ρ be a valuation for M and let $w \in W$. To show that (cnc) is valid in M , we suppose that $M, w \Vdash_{\rho} (\ell * j) \wedge (\ell * k)$, and require to show that $M, w \Vdash_{\rho} \@_j k$, i.e., that $\rho(j) = \rho(k)$. That $M, w \Vdash_{\rho} (\ell * j) \wedge (\ell * k)$ straightforwardly means that $w \in (\rho(\ell) \circ \rho(j)) \cap (\rho(\ell) \circ \rho(k))$. By cancellativity, we thus immediately get that $\rho(j) = \rho(k)$ as required.

(\Rightarrow) Assume that (cnc) is valid in M , and suppose $w' \in (w \circ w_1) \cap (w \circ w_2)$. We require to show that $w_1 = w_2$. Since (cnc) is valid in M , we have $M, w \Vdash_{\rho} (\text{cnc})$ for all $w \in W$ and hybrid valuations ρ . We define a hybrid valuation ρ for M as follows:

$$\rho(\ell) = w \quad \rho(j) = w_1 \quad \rho(k) = w_2$$

Then by assumption $M, w' \Vdash_{\rho} (\ell * j) \wedge (\ell * k)$. By validity of (cnc), we deduce that $M, w' \Vdash_{\rho} \@_j k$, hence that $\rho(j) = \rho(k)$ hence by construction that $w_1 = w_2$.

Single unit. (\Leftarrow) Assume $E = \{e\}$, let ρ be a valuation for M and let $w \in W$. We show (su) is valid. Supposing that

$M, w \Vdash_{\rho} \@_{\ell_1} I \wedge \@_{\ell_2} I$, we have that $\rho(\ell_1), \rho(\ell_2) \in E$. Thus $\rho(\ell_1) = \rho(\ell_2) = e$, and so we easily have $M, w \Vdash_{\rho} \@_{\ell_1} \ell_2$ as required.

(\Rightarrow) Assume (su) is valid and let $e, e' \in E$. We need to show $e = e'$. Define a valuation ρ for M by

$$\rho(\ell_1) = e \quad \rho(\ell_2) = e'$$

Thus, to show $e = e'$, it suffices to show $M, w \Vdash_{\rho} \@_{\ell_1} \ell_2$ (for any $w \in W$). Since (su) is valid, it suffices to show that $M, w \Vdash_{\rho} \@_{\ell_1} I \wedge \@_{\ell_2} I$. This follows immediately from our construction of ρ .

Disjointness. (\Leftarrow) Assume the disjointness property, let ρ be a valuation for M and let $w \in W$. To show (dis) is valid, we suppose that $M, w \Vdash_{\rho} \ell * \ell$ and require to show that $M, w \Vdash_{\rho} I \wedge \ell$, i.e., that $w = \rho(\ell)$ and $w \in E$. That $M, w \Vdash_{\rho} \ell * \ell$ means that $w \in \rho(\ell) \circ \rho(\ell)$. In particular, $\rho(\ell) \circ \rho(\ell) \neq \emptyset$ hence, by disjointness, $\rho(\ell) \in E$. We thus get $w \in \rho(\ell) \circ \rho(\ell) = \{\rho(\ell)\}$ hence $w = \rho(\ell)$ and also $w \in E$ as required.

(\Rightarrow) Assume that (dis) is valid in M , and suppose that $w \circ w \neq \emptyset$. We require to show that $w \in E$. Let $w' \in w \circ w$ and ρ be such that $\rho(\ell) = w$. By construction, we have that $M, w' \Vdash_{\rho} \ell * \ell$, hence by validity of (dis) we get $M, w' \Vdash_{\rho} I \wedge \ell$, i.e., $w' = \rho(\ell) = w$ and $w' \in E$, hence $w \in E$ as required. \square

Corollary 4.4. *Any separation theory not including the cross-split property is HyBBI-definable.*

Proof. Follows by taking as the defining formula the conjunction of the relevant formulas from Theorem 4.3 defining the properties of the theory. \square

As is also the case for BBI (see Conjecture 3.14), we do not currently know the status of definability of the cross-split property in HyBBI, but we suspect that it is not definable even with the extra power of nominals. This is because a straightforward translation of the property into HyBBI would require some way of binding or existentially quantifying nominals, which is not provided by pure nominals or the $\@_{\ell}$ modality. In Section 7 we add a binder to HyBBI, which enables us to express cross-split.

5. An axiomatic proof system for HyBBI

Here, we present a Hilbert-style axiomatic proof system for HyBBI, and show that it is sound with respect to validity in BBI-models; we examine questions of completeness in Section 6.

Definition 5.1. We define $\mathbf{K}_{\text{HyBBI}}$ to be the proof system obtained by extending the proof system \mathbf{K}_{BBI} (see Definition 2.4) with the axioms and rules for nominals and $\@$ given in Figure 1.

Our $\mathbf{K}_{\text{HyBBI}}$ is based on the proof system for basic hybrid logic in [1]. The axioms and rules have been chosen to make the subsequent completeness proof as simple as possible, rather than for minimality.

Proposition 5.2 (Soundness). *Any $\mathbf{K}_{\text{HyBBI}}$ -provable formula is valid.*

Proof. Let $M = \langle W, \circ, E \rangle$ be a BBI-model. Then, assuming A is $\mathbf{K}_{\text{HyBBI}}$ -provable, we must show that A is valid in M . It suffices to show that all axioms of $\mathbf{K}_{\text{HyBBI}}$ are valid and that validity is preserved by every proof rule of $\mathbf{K}_{\text{HyBBI}}$. This is a straightforward verification for all the rules and axioms except the two ‘‘bridge’’ axioms and the two ‘‘paste’’ rules. We just show the cases

$(K_{\textcircled{a}})$ $\textcircled{a}_\ell(A \rightarrow B) \vdash \textcircled{a}_\ell A \rightarrow \textcircled{a}_\ell B$ (Self-dual) $\textcircled{a}_\ell A \vdash \neg \textcircled{a}_\ell \neg A$ and $\neg \textcircled{a}_\ell \neg A \vdash \textcircled{a}_\ell A$ $(\textcircled{a}\text{-intro})$ $\ell \wedge A \vdash \textcircled{a}_\ell A$ (Refl) $\vdash \textcircled{a}_\ell \ell$ (Sym) $\textcircled{a}_\ell k \vdash \textcircled{a}_k \ell$	(Nom) $\textcircled{a}_\ell k \wedge \textcircled{a}_k A \vdash \textcircled{a}_\ell A$ (Agree) $\textcircled{a}_k \textcircled{a}_\ell A \vdash \textcircled{a}_\ell A$ and $\textcircled{a}_\ell A \vdash \textcircled{a}_k \textcircled{a}_\ell A$ $(\text{Bridge } *)$ $\textcircled{a}_\ell(k * k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B \vdash \textcircled{a}_\ell(A * B)$ $(\text{Bridge } \rightarrow)$ $\textcircled{a}_\ell \neg(k \rightarrow \neg k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B \vdash \textcircled{a}_\ell \neg(A \rightarrow \neg B)$		
$\frac{\vdash A}{\vdash A[\theta]} (\text{Subst})$	$\frac{\vdash A}{\vdash \textcircled{a}_\ell A} (\text{Gen})$	$\frac{\textcircled{a}_\ell(k * k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B \vdash C}{\textcircled{a}_\ell(A * B) \vdash C} \quad k, k' \text{ not in } A, B, C \text{ or } \{\ell\}$	$\frac{\textcircled{a}_\ell \neg(k \rightarrow \neg k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B \vdash C}{\textcircled{a}_\ell \neg(A \rightarrow \neg B) \vdash C} \quad k, k' \text{ not in } A, B, C \text{ or } \{\ell\}$
$\frac{\ell \vdash A}{\vdash A} \quad \ell \text{ not in } A \text{ (Name)}$	$\frac{\textcircled{a}_\ell \neg(k \rightarrow \neg k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B \vdash C}{\textcircled{a}_\ell \neg(A \rightarrow \neg B) \vdash C} \quad k, k' \text{ not in } A, B, C \text{ or } \{\ell\}$	$\frac{\textcircled{a}_\ell(k * k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B \vdash C}{\textcircled{a}_\ell(A * B) \vdash C} \quad k, k' \text{ not in } A, B, C \text{ or } \{\ell\}$	$\frac{\textcircled{a}_\ell \neg(k \rightarrow \neg k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B \vdash C}{\textcircled{a}_\ell \neg(A \rightarrow \neg B) \vdash C} \quad k, k' \text{ not in } A, B, C \text{ or } \{\ell\}$

Figure 1. Rules and axioms for nominals in $\mathbf{K}_{\text{HyBBI}}$. Note that θ in the the rule (Subst) is a substitution of nominals for nominals.

of (Bridge $*$) and (Paste \rightarrow) here, as the others are similar.

Case (Bridge $$).* Let ρ be a valuation for M and let $w \in W$. Suppose $M, w \models_\rho \textcircled{a}_\ell(k * k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B$. Then we have $M, \rho(\ell) \models_\rho k * k'$ and $M, \rho(k) \models_\rho A$ and $M, \rho(k') \models_\rho B$. The first of these means that $\rho(\ell) \in w_1 \circ w_2$ and $M, w_1 \models_\rho k$ and $M, w_2 \models_\rho k'$. That is, $\rho(\ell) \in \rho(k) \circ \rho(k')$. Thus $M, \rho(\ell) \models_\rho A * B$, i.e. $M, w \models_\rho \textcircled{a}_\ell(A * B)$ as required.

Case (Paste \rightarrow). Let ρ be a valuation for M and let $w \in W$. Supposing the premise of the rule is valid in M and $M, w \models_\rho \textcircled{a}_\ell \neg(A \rightarrow \neg B)$, we have to show $M, w \models_\rho C$. We have $M, \rho(\ell) \models_\rho \neg(A \rightarrow \neg B)$ which means that there exist $w', w'' \in W$ such that $w'' \in \rho(\ell) \circ w'$ and $M, w' \models_\rho A$ and $M, w'' \models_\rho B$. Now define the valuation $\rho' = \rho[k \mapsto w', k' \mapsto w'']$, where k and k' are the fresh nominals appearing in the premise of the rule. By construction, and using the fact that ρ and ρ' agree except possibly on the fresh nominals k, k' , we have $\rho'(k') \in \rho'(k) \circ \rho'(k)$ and $M, \rho'(k) \models_{\rho'} A$ and $M, \rho'(k') \models_{\rho'} B$. The first of these gives us $M, \rho'(\ell) \models_{\rho'} \neg(k \rightarrow \neg k')$. Putting everything together, we obtain

$$M, w \models_{\rho'} \textcircled{a}_\ell \neg(k \rightarrow \neg k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} B$$

Since the premise of the rule is assumed valid, we obtain $M, w \models_{\rho'} C$. Again, since ρ and ρ' agree except on k, k' , which do not appear in C , we thus obtain $M, w \models_\rho C$ as required. \square

The following example illustrates how the hybrid axioms and rules are used in practice.

Example 5.3. The HyBBI-formula $\top * (I \wedge \neg(\ell \rightarrow \neg A)) \vdash \textcircled{a}_\ell A$ is provable in HyBBI.

Proof. Intuitively, the LHS of the formula says that one may find an empty sub-state to the current state which one can compose with $\rho(\ell)$ to satisfy A . This should imply that $\rho(\ell)$ itself satisfies A .

First, we show that the following formula is provable:

$$A * \textcircled{a}_\ell B \vdash \textcircled{a}_\ell B \quad (1)$$

Let j, k, k' be fresh nominals not occurring in A, B or $\{\ell\}$. We have $\textcircled{a}_{k'} \textcircled{a}_\ell B \vdash \textcircled{a}_\ell B$ an instance of (Agree). By weakening for \wedge , we thus obtain

$$\textcircled{a}_j(k * k') \wedge \textcircled{a}_k A \wedge \textcircled{a}_{k'} \textcircled{a}_\ell B \vdash \textcircled{a}_\ell B$$

Using the fact that k, k' are fresh, we can apply the rule (Paste $*$) to derive $\textcircled{a}_j(A * \textcircled{a}_\ell B) \vdash \textcircled{a}_\ell B$. Since the formula

$$j \wedge (A * \textcircled{a}_\ell B) \vdash \textcircled{a}_j(A * \textcircled{a}_\ell B)$$

is an instance of (\textcircled{a} -intro), we obtain $j \wedge (A * \textcircled{a}_\ell B) \vdash \textcircled{a}_\ell B$ by transitivity, and thus easily we have $j \vdash (A * \textcircled{a}_\ell B) \rightarrow \textcircled{a}_\ell B$. Since

j is fresh, we obtain $A * \textcircled{a}_\ell B \vdash \textcircled{a}_\ell B$ as required by applying (Name).

Next, we show that the following formula is provable:

$$I \wedge \neg(\ell \rightarrow \neg A) \vdash \textcircled{a}_\ell A \quad (2)$$

We have $\ell \wedge A \vdash \textcircled{a}_\ell A$ an instance of (\textcircled{a} -intro), whence by contraposition and use of (Self-dual) we obtain $\ell \wedge \textcircled{a}_\ell \neg A \vdash \neg A$. Now, since $\ell * \textcircled{a}_\ell \neg A \vdash \textcircled{a}_\ell \neg A$ is provable as an instance of (1) above, we obtain $\ell \wedge (\ell * \textcircled{a}_\ell \neg A) \vdash \neg A$. By straightforward manipulations of plain BBI we can prove

$$(I \wedge \textcircled{a}_\ell \neg A) * \ell \vdash \ell \wedge (\textcircled{a}_\ell \neg A * \ell)$$

Thus, by transitivity, we obtain $(I \wedge \textcircled{a}_\ell \neg A) * \ell \vdash \neg A$. This rearranges to $I \wedge \textcircled{a}_\ell \neg A \vdash \ell \rightarrow \neg A$ and then to $I \wedge \neg(\ell \rightarrow \neg A) \vdash \neg \textcircled{a}_\ell \neg A$, which yields the required (2) by using (Self-dual).

Now we can supply the required derivation of the formula in the proposition. We can derive $\top * (I \wedge \neg(\ell \rightarrow \neg A)) \vdash \top * \textcircled{a}_\ell A$ using (2) and the BBI proof rules for $*$. As $\top * \textcircled{a}_\ell A \vdash \textcircled{a}_\ell A$ is an instance of (1), we have $\top * (I \wedge \neg(\ell \rightarrow \neg A)) \vdash \textcircled{a}_\ell A$ by transitivity as required. \square

Interestingly, the converse of the formula in Example 5.3, that is $\textcircled{a}_\ell A \vdash \top * (I \wedge \neg(\ell \rightarrow \neg A))$, is not generally valid, but is valid in all single-unit models (and thus in such models \textcircled{a}_ℓ is definable already using plain nominals). Roughly speaking, the RHS of the formula says that given any starting world $w \in W$ we have $w = w \circ e$ for some $e \in E$ such that $\rho(\ell) \circ e$ is defined. This is true in models with a single unit, but might fail in models with multiple units. In general, the extra expressivity of the \textcircled{a}_ℓ modality beyond that yielded by pure nominals is that it enables us to talk about worlds not accessible from the current world via the $I, *$ and \rightarrow modalities. (In single-unit models, there are no such worlds.)

6. Completeness for pure extensions of $\mathbf{K}_{\text{HyBBI}}$

In this section, we show a parametric completeness result: any extension of $\mathbf{K}_{\text{HyBBI}}$ with a set of pure axioms Ax is complete with respect to the class of BBI-models satisfying Ax. In particular, we can obtain complete proof systems for many separation theories simply by adding the axioms defining the theory to $\mathbf{K}_{\text{HyBBI}}$.

We follow the basic structure of the corresponding completeness proof for normal hybrid logic in [1], which shows that any consistent set of formulas has a model based upon “named” maximal consistent sets. Compared to this proof, we encounter two additional difficulties. First, we have to work with the residuated binary connectives $*$ and \rightarrow , as opposed to a single diamond modality and its De Morgan dual. Second, we have to show that the model we construct is a BBI-model, as opposed to an unrestricted frame.

Definition 6.1 (Consistent set). Let \mathcal{K} be any proof system. A set Γ of formulas is said to be \mathcal{K} -inconsistent if there are formulas $A_1, \dots, A_n \in \Gamma$ such that $A_1 \wedge \dots \wedge A_n \vdash \perp$ is provable in \mathcal{K} . Otherwise Γ is called \mathcal{K} -consistent.

Definition 6.2 (Maximal consistent set). Let \mathcal{K} be any proof system. A set Γ of formulas is *maximal \mathcal{K} -consistent* (and we call Γ a \mathcal{K} -MCS) if Γ is \mathcal{K} -consistent and any $\Delta \supset \Gamma$ is \mathcal{K} -inconsistent.

In the rest of this section, whenever we talk about MCSs, consistency and provability, we *always* mean with reference to an arbitrary extension $\mathbf{K}_{\text{HyBBI}} + \text{Ax}$ of $\mathbf{K}_{\text{HyBBI}}$ with a finite set Ax of axioms expressed as pure formulas.

We begin by recalling some basic facts about MCSs.

Lemma 6.3. *For any MCS Γ and formulas A, B , we have*

1. if $A \vdash B$ is provable and $A \in \Gamma$ then $B \in \Gamma$;
2. $\top \in \Gamma$ and $\perp \notin \Gamma$;
3. either $A \in \Gamma$ or $\neg A \in \Gamma$;
4. $A \wedge B \in \Gamma$ iff $A, B \in \Gamma$;
5. $A \vee B \in \Gamma$ iff $A \in \Gamma$ or $B \in \Gamma$.

Proof. Standard in all cases. \square

In the following, we do not refer explicitly to uses of Lemma 6.3, as we use it so frequently.

Definition 6.4 (Named / pasted MCS). An MCS Γ is said to be *named* if there is at least one nominal $\ell \in \Gamma$; any such ℓ is called a *name* for Γ .

Γ is said to be *pasted* if

- $\@_{\ell}(A * B) \in \Gamma$ implies $\@_{\ell}(\ell_1 * \ell_2) \wedge \@_{\ell_1}A \wedge \@_{\ell_2}B \in \Gamma$ for some ℓ_1, ℓ_2 , and
- $\@_{\ell}\neg(A * \neg B) \in \Gamma$ implies $\@_{\ell}\neg(\ell_1 * \neg \ell_2) \wedge \@_{\ell_1}A \wedge \@_{\ell_2}B \in \Gamma$ for some ℓ_1, ℓ_2 .

Lemma 6.5 (Extended Lindenbaum Lemma). *Let \mathcal{N}' be a countably infinite set of nominals disjoint from \mathcal{N} . If Δ is a consistent set of formulas then there is a named, pasted MCS Δ^+ (of formulas in the extended nominal language of $\mathcal{N} \cup \mathcal{N}'$) such that $\Delta \subseteq \Delta^+$.*

Proof. Let k_0, k_1, k_2, \dots be an enumeration of \mathcal{N}' , and let B_1, B_2, B_3, \dots be an enumeration of all formulas in the extended language given by $\mathcal{N} \cup \mathcal{N}'$. Given a consistent set Δ of formulas, we define a sequence $(\Delta_i)_{i \geq 0}$ of sets of formulas as follows:

- $\Delta_0 \stackrel{\text{def}}{=} \Delta \cup \{k_0\}$;
- if $\Delta_i \cup \{B_i\}$ is inconsistent then $\Delta_{i+1} \stackrel{\text{def}}{=} \Delta_i$;
- if $\Delta_i \cup \{B_i\}$ is consistent and the formula B_i is not of the form $\@_{\ell}(A * B)$ or $\@_{\ell}\neg(A * \neg B)$, then $\Delta_{i+1} \stackrel{\text{def}}{=} \Delta_i \cup \{B_i\}$;
- if $\Delta_i \cup \{B_i\}$ is consistent and $B_i = \@_{\ell}(A * B)$ then

$$\Delta_{i+1} \stackrel{\text{def}}{=} \Delta_i \cup \{B_i\} \cup \{\@_{\ell}(k * k') \wedge \@_k A \wedge \@_{k'} B\}$$

where k, k' are fresh nominals from our enumeration of \mathcal{N}' ;

- if $\Delta_i \cup \{B_i\}$ is consistent and $B_i = \@_{\ell}\neg(A * \neg B)$ then

$$\Delta_{i+1} \stackrel{\text{def}}{=} \Delta_i \cup \{B_i\} \cup \{\@_{\ell}\neg(k * \neg k') \wedge \@_k A \wedge \@_{k'} B\}$$

where k, k' in the last two clauses are fresh nominals from our enumeration of \mathcal{N}' . We claim that $\Delta^+ \stackrel{\text{def}}{=} \bigcup_{i \geq 0} \Delta_i$ is a named, pasted MCS.

First, to see that Δ^+ is consistent, it suffices to show that Δ_i is consistent for all i . We proceed by induction on i . In the case $i = 0$, we must show that $\Delta \cup \{k_0\}$ is consistent. If not, then there are formulas $A_1, \dots, A_n \in \Delta$ such that, writing $A = \bigwedge_{1 \leq i \leq n} A_i$, we have $A \wedge k_0 \vdash \perp$ provable. Thus $k_0 \vdash \neg A$ is provable, whence by the rule (Name) we have $\vdash \neg A$ provable and thus $A \vdash \perp$ provable,

contradicting the consistency of Δ . Now, assuming that Δ_i is consistent, we must show that Δ_{i+1} is consistent. This is immediate by induction hypothesis except in the case that $B_i = \@_{\ell}(A * B)$ or $B_i = \@_{\ell}\neg(A * \neg B)$. We show the case $B_i = \@_{\ell}(A * B)$. In this case, assume for contradiction that there are $A_1, \dots, A_n \in \Delta_i$ such that, writing $A = \bigwedge_{1 \leq i \leq n} A_i$, the following is provable:

$$A \wedge \@_{\ell}(A * B) \wedge \@_{\ell}(k * k') \wedge \@_k A \wedge \@_{k'} B \vdash \perp$$

Thus we can also prove

$$\@_{\ell}(k * k') \wedge \@_k A \wedge \@_{k'} B \vdash \neg A \vee \neg \@_{\ell}(A * B)$$

Since k, k' are fresh nominals by construction, we obtain by applying (Paste *):

$$\@_{\ell}(A * B) \vdash \neg A \vee \neg \@_{\ell}(A * B)$$

Thus we obtain $A \wedge \@_{\ell}(A * B) \vdash \perp$, contradicting the assumed consistency of $\Delta_i \cup \{B_i\}$. The case $B_i = \neg(A * \neg B)$ is similar, using the rule (Paste \neg).

Next, we must show that Δ^+ is maximal. Suppose that for some formula A , we have $\Delta^+ \cup \{A\}$ consistent but $A \notin \Delta^+$. Note that A appears in our enumeration as B_i say, so by construction it must be that $\Delta_i \cup \{A\}$ is inconsistent (otherwise $A \in \Delta_{i+1} \subseteq \Delta^+$). But then $\Delta^+ \cup \{A\}$ is inconsistent, contradiction.

Next, to see that Δ^+ is named, observe that $k_0 \in \Delta_0 \subseteq \Delta^+$ by construction.

Finally, we show Δ^+ is pasted. First, suppose $\@_{\ell}(A * B) \in \Delta^+$. Note that $\@_{\ell}(A * B)$ appears as some B_i in our enumeration. Now every finite subset of an MCS is consistent, so $\Delta_i \cup \{B_i\}$ must be consistent. Thus, by construction, we have

$$\@_{\ell}(k * k') \wedge \@_k A \wedge \@_{k'} B \in \Delta_{i+1} \subseteq \Delta^+$$

as required. For similar reasons, whenever $\@_{\ell}\neg(A * \neg B) \in \Delta^+$ we have $\@_{\ell}\neg(k * \neg k') \wedge \@_k A \wedge \@_{k'} B \in \Delta^+$. This completes the proof. \square

In the following, we define a *named set yielded* by Γ to be any set of formulas $\{A \mid \@_{\ell}A \in \Gamma\}$ for some nominal ℓ .

Lemma 6.6. *Let Γ be an MCS, and let $\Delta_{\ell} \stackrel{\text{def}}{=} \{A \mid \@_{\ell}A \in \Gamma\}$ be a named set yielded by Γ for each nominal ℓ . Then the following hold for all nominals ℓ, k :*

1. Δ_{ℓ} is an MCS containing ℓ ;
2. if $\ell \in \Delta_k$ then $\Delta_k = \Delta_{\ell}$;
3. $\@_{\ell}A \in \Delta_k$ iff $\@_{\ell}A \in \Gamma$;
4. if ℓ is a name for Γ then $\Gamma = \Delta_{\ell}$.

Proof. The proof of the analogous result for normal hybrid logic, stated as Lemma 7.24 in [1], also suffices for our setting. \square

Definition 6.7. A BBI-frame $\langle W, \circ, E \rangle$ is said to be *named* by the hybrid valuation ρ if for all $w \in W$ there is an $\ell \in \mathcal{N}$ with $\rho(\ell) = w$.

Definition 6.8. Let Γ be a named, pasted MCS. Then the *named model yielded* by Γ is defined as $M^{\Gamma} \stackrel{\text{def}}{=} \langle W^{\Gamma}, \circ^{\Gamma}, E^{\Gamma} \rangle$, where:

1. W^{Γ} is the set of all named sets yielded by Γ ;
2. $\Delta_1 \circ^{\Gamma} \Delta_2 \stackrel{\text{def}}{=} \{\Delta \mid A_1 \in \Delta_1, A_2 \in \Delta_2 \text{ implies } A_1 * A_2 \in \Delta\}$;
3. $E^{\Gamma} \stackrel{\text{def}}{=} \{\Delta \mid \text{I} \in \Delta\}$.

The *canonical valuation* ρ^{Γ} for M^{Γ} is defined by

$$\rho^{\Gamma}(P) \stackrel{\text{def}}{=} \{\Delta \mid P \in \Delta\} \quad P \text{ a proposition}$$

$$\rho^{\Gamma}(\ell) \stackrel{\text{def}}{=} \{A \mid \@_{\ell}A \in \Gamma\} \quad \ell \text{ a nominal}$$

We observe that M^Γ is indeed a BBI-frame named by ρ^Γ : for any $\Delta \in W^\Gamma$ we have $\Delta = \{A \mid @_\ell A \in \Gamma\}$ for some ℓ , whence by definition $\rho^\Gamma(\ell) = \Delta$. We will show that it is also a BBI-model in Lemma 6.13, and one such that $M^\Gamma, \Delta \models_{\rho^\Gamma} A$ if and only if $A \in \Delta$ for all Δ, A in Lemma 6.12. In the meantime, let us present a number of intermediary lemmas.

Lemma 6.9 (Existence Lemma for $*$). *For any $\Delta \in W^\Gamma$, if $A_1 * A_2 \in \Delta$ then there exist $\Delta_1, \Delta_2 \in W^\Gamma$ such that $\Delta \in \Delta_1 \circ^\Gamma \Delta_2$ and $A_1 \in \Delta_1, A_2 \in \Delta_2$.*

Proof. Let $A_1 * A_2 \in \Delta$. We have $\Delta = \{A \mid @_\ell A \in \Gamma\}$ for some nominal ℓ . Thus $@_\ell(A_1 * A_2) \in \Gamma$. As Γ is pasted, we have nominals ℓ_1, ℓ_2 such that $@_\ell(\ell_1 * \ell_2) \wedge @_{\ell_1} A_1 \wedge @_{\ell_2} A_2 \in \Gamma$. Thus $A_1 \in \Delta_1$ and $A_2 \in \Delta_2$, where $\Delta_1 = \{A \mid @_{\ell_1} A \in \Gamma\}$ and $\Delta_2 = \{A \mid @_{\ell_2} A \in \Gamma\}$ are named sets yielded by Γ .

It just remains to show that $\Delta \in \Delta_1 \circ^\Gamma \Delta_2$. Let $B_1 \in \Delta_1, B_2 \in \Delta_2$. By definition, $@_{\ell_1} B_1 \in \Gamma$ and $@_{\ell_2} B_2 \in \Gamma$. As MCSs are closed under provability and conjunction, we have $@_\ell(\ell_1 * \ell_2) \wedge @_{\ell_1} B_1 \wedge @_{\ell_2} B_2 \in \Gamma$. Thus, using the rule (Bridge $*$), we have $@_\ell(B_1 * B_2) \in \Gamma$. Thus $B_1 * B_2 \in \Delta$ as required. \square

Lemma 6.10. $\Delta \in \Delta_1 \circ^\Gamma \Delta_2$ if and only if for all formulas A and $B, A \in \Delta_2$ and $B \in \Delta$ implies $\neg(A \multimap \neg B) \in \Delta_1$.

Proof. We show each direction of the bimplication separately, making use of the fact that $\Delta, \Delta_1, \Delta_2$ are MCSs by part 1 of Lemma 6.6.

(\Leftarrow) Let $A_1 \in \Delta_1, A_2 \in \Delta_2$ and suppose for contradiction that $A_1 * A_2 \notin \Delta$. As Δ is an MCS, $\neg(A_1 * A_2) \in \Delta$. By assumption, $\neg(A_2 \multimap \neg(A_1 * A_2)) \in \Delta_1$. As Δ_1 is an MCS, we have $A_1 \wedge \neg(A_2 \multimap (A_1 * A_2)) \in \Delta_1$. But $A_1 \wedge \neg(A_2 \multimap (A_1 * A_2)) \vdash \perp$ is provable (since $A_1 \vdash A_2 \multimap (A_1 * A_2)$ is provable). This contradicts the consistency of Δ_1 . Hence $A_1 * A_2 \in \Delta$ as required.

(\Rightarrow) Let $A \in \Delta_2, B \in \Delta$ and suppose for contradiction that $\neg(A \multimap \neg B) \notin \Delta_1$. As Δ_1 is an MCS, we have $A \multimap \neg B \in \Delta_1$, so by the main assumption $(A \multimap \neg B) * A \in \Delta$. As Δ is an MCS and $(A \multimap \neg B) * A \vdash \neg B$ is provable, $\neg B \in \Delta$. This contradicts the consistency of Δ , so $\neg(A \multimap \neg B) \in \Delta_1$ as required. \square

Lemma 6.11 (Existence Lemma for \multimap). *For any $\Delta \in W^\Gamma$, if $\neg(A_1 \multimap \neg A_2) \in \Delta$ then there exist $\Delta', \Delta'' \in W^\Gamma$ such that $\Delta'' \in \Delta \circ^\Gamma \Delta'$ and $A_1 \in \Delta', A_2 \in \Delta''$.*

Proof. Assume $\neg(A_1 \multimap \neg A_2) \in \Delta$. We have $\Delta = \{A \mid @_\ell A \in \Gamma\}$ for some nominal ℓ . Thus $@_\ell \neg(A_1 \multimap \neg A_2) \in \Gamma$. Since Γ is pasted, we have $@_\ell \neg(\ell_1 \multimap \neg \ell_2) \wedge @_{\ell_1} A_1 \wedge @_{\ell_2} A_2 \in \Gamma$ for some ℓ_1, ℓ_2 . Thus $A_1 \in \Delta'$ and $A_2 \in \Delta''$, where $\Delta' = \{A \mid @_{\ell_1} A \in \Gamma\}$ and $\Delta'' = \{A \mid @_{\ell_2} A \in \Gamma\}$ are named sets yielded by Γ .

It just remains to show that $\Delta'' \in \Delta \circ^\Gamma \Delta'$. According to Lemma 6.10, it suffices to show that $A \in \Delta'$ and $B \in \Delta''$ implies $\neg(A \multimap \neg B) \in \Delta$. Supposing $A \in \Delta', B \in \Delta''$, we have $@_{\ell_1} A \in \Gamma$ and $@_{\ell_2} B \in \Gamma$. Thus we obtain $@_\ell \neg(\ell_1 \multimap \neg \ell_2) \wedge @_{\ell_1} A \wedge @_{\ell_2} B \in \Gamma$. Since Γ is an MCS it is closed under the rule (Bridge \multimap), whence $@_\ell \neg(A \multimap \neg B) \in \Gamma$. Thus $\neg(A \multimap \neg B) \in \Delta$ as required. \square

Lemma 6.12 (Truth Lemma). *For any HyBBI-formula A and $\Delta \in W^\Gamma$, we have $M^\Gamma, \Delta \models_{\rho^\Gamma} A$ if and only if $A \in \Delta$.*

Proof. By structural induction on A . Throughout, we make use of the fact that any named set yielded by Γ is also an MCS, which is guaranteed by part 1 of Lemma 6.6.

Case $A = P$. Using the definition of ρ^Γ , we have as required:

$$M^\Gamma, \Delta \models_{\rho^\Gamma} P \Leftrightarrow \Delta \in \rho^\Gamma(P) \Leftrightarrow P \in \Delta$$

Case $A = \ell$. Using the definition of ρ^Γ , we have

$$M^\Gamma, \Delta \models_{\rho^\Gamma} \ell \Leftrightarrow \Delta = \rho^\Gamma(\ell) \Leftrightarrow \Delta = \{A \mid @_\ell A \in \Gamma\}$$

Now, going from left to right, we have $\Delta = \{A \mid @_\ell A \in \Gamma\}$ and thus $\ell \in \Delta$ by part 1 of Lemma 6.6. Conversely, assuming $\ell \in \Delta$, we have that $\Delta = \{A \mid @_\ell A \in \Gamma\}$ for some ℓ , and by part 2 of Lemma 6.6 we obtain $\Delta = \{A \mid @_\ell A \in \Gamma\}$ as required.

Case $A = \top$. As \top is in any MCS, we have $\top \in \Delta$, so as required

$$M^\Gamma, \Delta \models_{\rho^\Gamma} \top \Leftrightarrow \text{always} \Leftrightarrow \top \in \Delta$$

Case $A = \perp$. As \perp cannot be in any MCS, we have $\perp \notin \Delta$, so as required

$$M^\Gamma, \Delta \models_{\rho^\Gamma} \perp \Leftrightarrow \text{never} \Leftrightarrow \perp \in \Delta$$

Case $A = \neg A'$. Using the induction hypothesis for A' , we have

$$M^\Gamma, \Delta \models_{\rho^\Gamma} \neg A' \Leftrightarrow M^\Gamma, \Delta \not\models_{\rho^\Gamma} A' \Leftrightarrow A' \notin \Delta$$

Since Δ is an MCS, we have $A' \notin \Delta$ if and only if $\neg A' \in \Delta$, which completes the case.

Case $A = A_1 \wedge A_2$. Using the induction hypothesis for A_1 and A_2 , we have:

$$\begin{aligned} M^\Gamma, \Delta \models_{\rho^\Gamma} A_1 \wedge A_2 \\ \Leftrightarrow M^\Gamma, \Delta \models_{\rho^\Gamma} A_1 \text{ and } M^\Gamma, \Delta \models_{\rho^\Gamma} A_2 \\ \Leftrightarrow A_1 \in \Delta \text{ and } A_2 \in \Delta \end{aligned}$$

To complete the case, we observe that as Δ is an MCS it is the case that $A_1, A_2 \in \Delta$ if and only if $A_1 \wedge A_2 \in \Delta$.

Cases $A = A_1 \vee A_2, A = A_1 \rightarrow A_2$. Similar to the previous case.

Case $A = I$. Using the definition of E^Γ , we easily have as required:

$$M^\Gamma, \Delta \models_{\rho^\Gamma} I \Leftrightarrow \Delta \in E^c \Leftrightarrow I \in \Delta$$

*Case $A = A_1 * A_2$.* Using the induction hypothesis for A_1 and A_2 , we have:

$$\begin{aligned} M^\Gamma, \Delta \models_{\rho^\Gamma} A_1 * A_2 \\ \Leftrightarrow \Delta \in \Delta_1 \circ^\Gamma \Delta_2 \text{ and } M^\Gamma, \Delta_1 \models_{\rho^\Gamma} A_1 \text{ and } M^\Gamma, \Delta_2 \models_{\rho^\Gamma} A_2 \\ \Leftrightarrow \Delta \in \Delta_1 \circ^\Gamma \Delta_2 \text{ and } A_1 \in \Delta_1 \text{ and } A_2 \in \Delta_2 \end{aligned}$$

Thus, going from left to right, we immediately get $A_1 * A_2 \in \Delta$ as required from the definition of \circ^Γ .

Going from right to left, we have $A_1 * A_2 \in \Delta$ and must construct the required named sets Δ_1, Δ_2 yielded by Γ satisfying the statement above. This is precisely guaranteed by our Existence Lemma for $*$ (Lemma 6.9).

Case $A = A_1 \multimap A_2$. Using the induction hypothesis for A_1 and A_2 , we have:

$$\begin{aligned} M^\Gamma, \Delta \models_{\rho^\Gamma} A_1 \multimap A_2 \\ \Leftrightarrow \forall \Delta', \Delta''. \Delta'' \in \Delta \circ^\Gamma \Delta' \text{ and } M^\Gamma, \Delta' \models_{\rho^\Gamma} A_1 \text{ implies} \\ M^\Gamma, \Delta'' \models_{\rho^\Gamma} A_2 \\ \Leftrightarrow \forall \Delta', \Delta''. \Delta'' \in \Delta \circ^\Gamma \Delta' \text{ and } A_1 \in \Delta' \text{ implies } A_2 \in \Delta'' \end{aligned}$$

Thus, going from right to left, assume that $A_1 \multimap A_2 \in \Delta, A_1 \in \Delta'$ and $\Delta'' \in \Delta \circ^\Gamma \Delta'$. By the definition of \circ^Γ , we have $(A_1 \multimap A_2) * A_1 \in \Delta''$, whence we obtain $A_2 \in \Delta''$ as required by modus ponens.

Going from left to right, however, we must show that $A_1 \multimap A_2 \in \Delta$ given the above implication. We show the contrapositive. Assume that $A_1 \multimap A_2 \notin \Delta$, which since Δ is an MCS means

that $\neg(A_1 \multimap A_2) \in \Delta$. We must construct named sets Δ', Δ'' yielded by Γ , with $\Delta'' \in \Delta \circ^\Gamma \Delta'$ and $A_1 \in \Delta'$ but $A_2 \notin \Delta''$, i.e. $\neg A_2 \in \Delta$. This is provided by our Existence Lemma for \multimap (Lemma 6.11).

Case $A = @_\ell B$. Using the induction hypothesis for B , we have:

$$\begin{aligned} M^\Gamma, \Delta \models_{\rho^\Gamma} @_\ell B &\Leftrightarrow M^\Gamma, \rho^\Gamma(\ell) \models_{\rho^\Gamma} B \\ &\Leftrightarrow B \in \rho^\Gamma(\ell) \\ &\Leftrightarrow B \in \{A \mid @_\ell A \in \Gamma\} \\ &\Leftrightarrow @_\ell B \in \Gamma \end{aligned}$$

Now, using part 3 of Lemma 6.6, we have that $@_\ell B \in \Gamma$ if and only if $@_\ell B \in \Delta$ (since Δ is a named set yielded by Γ). This completes the case, and the proof. \square

Lemma 6.13. *Let $M^\Gamma = \langle W^\Gamma, \circ^\Gamma, E^\Gamma \rangle$ be the named model yielded by the named, pasted MCS Γ . Then M^Γ is a BBI-model.*

Proof. To show that M^Γ is a BBI-model, we have to show that it satisfies the axioms given in Definition 2.2.

Commutativity. It suffices to show that $\Delta_1 \circ^\Gamma \Delta_2 \subseteq \Delta_2 \circ^\Gamma \Delta_1$. Let $\Delta \in \Delta_1 \circ^\Gamma \Delta_2$, and suppose $A_1 \in \Delta_1, A_2 \in \Delta_2$. To show $\Delta \in \Delta_2 \circ^\Gamma \Delta_1$, we have to show $A_2 * A_1 \in \Delta$. As $\Delta \in \Delta_1 \circ^\Gamma \Delta_2$, we have $A_1 * A_2 \in \Delta$. As MCSs are closed under modus ponens and $A_1 * A_2 \vdash A_2 * A_1$ is provable, we have $A_2 * A_1 \in \Delta$ as required.

Associativity. It suffices by commutativity to show $\Delta_1 \circ^\Gamma (\Delta_2 \circ^\Gamma \Delta_3) \subseteq (\Delta_1 \circ^\Gamma \Delta_2) \circ^\Gamma \Delta_3$. Assume that $\Delta \in \Delta_1 \circ^\Gamma (\Delta_2 \circ^\Gamma \Delta_3)$, which means that for some $\Delta' \in \Delta_2 \circ^\Gamma \Delta_3$ we have $\Delta \in \Delta_1 \circ^\Gamma \Delta'$. Using part 1 of Lemma 6.6, we have ℓ_1, ℓ_2, ℓ_3 such that $\ell_i \in \Delta_i$ for each $i \in \{1, 2, 3\}$. As $\Delta' \in \Delta_2 \circ^\Gamma \Delta_3$, we have $\ell_2 * \ell_3 \in \Delta'$. Thus, as $\Delta \in \Delta_1 \circ^\Gamma \Delta'$, we have $\ell_1 * (\ell_2 * \ell_3) \in \Delta$. By applying associativity, $\ell_1 * (\ell_2 * \ell_3) \vdash (\ell_1 * \ell_2) * \ell_3$ is provable, so $(\ell_1 * \ell_2) * \ell_3 \in \Delta$. By two applications of the Existence Lemma for $*$ (Lemma 6.9) we obtain named sets $\Sigma_1, \Sigma_2, \Sigma_3, \Delta'' \in W^\Gamma$ such that $\ell_i \in \Sigma_i$ for each $i \in \{1, 2, 3\}$, and $\Delta \in \Delta'' \circ^\Gamma \Sigma_3$ and $\Delta'' \in \Sigma_1 \circ^\Gamma \Sigma_2$. By part 2 of Lemma 6.6, $\Sigma_i = \Delta_i$ for each $i \in \{1, 2, 3\}$. Hence $\Delta \in (\Delta_1 \circ^\Gamma \Delta_2) \circ^\Gamma \Delta_3$ as required.

Unit law. We must show that $E^\Gamma \circ^\Gamma \Delta = \{\Delta\}$ for any $\Delta \in W^\Gamma$. First we show that $E^\Gamma \circ^\Gamma \Delta \subseteq \{\Delta\}$. Suppose $\Delta' \in E^\Gamma \circ^\Gamma \Delta$, i.e. there is a $\Delta_E \in E^\Gamma$ such that $\Delta' \in \Delta_E \circ^\Gamma \Delta$. We need to show $\Delta' = \Delta$. First suppose $A \in \Delta$, and note that $I \in \Delta_E$ by definition. By definition of \circ^Γ we have $I * A \in \Delta'$, and as $I * A \vdash A$ is provable we must have $A \in \Delta'$. Thus $\Delta' \supseteq \Delta$. To see that $\Delta' = \Delta$ as required, we just observe that if $\Delta' \supset \Delta$ then, as Δ' is consistent, Δ is not maximal, contradiction.

We still need to show that $\Delta \in E^\Gamma \circ^\Gamma \Delta$, i.e. that $\Delta \in \Delta_E \circ^\Gamma \Delta$ for some $\Delta_E \in E^\Gamma$. Using part 1 of Lemma 6.6, we have some $\ell \in \Delta$. Since $\ell \vdash I * \ell$ is provable, we have $I * \ell \in \Delta$. Using the Existence Lemma for $*$ (Lemma 6.9) we obtain named sets $\Delta_E, \Delta' \in W^\Gamma$ such that $\Delta \in \Delta_E \circ^\Gamma \Delta'$ and $I \in \Delta_E$ and $\ell \in \Delta'$. Thus $\Delta_E \in E^\Gamma$ and, by part 2 of Lemma 6.6, $\Delta' = \Delta$. This completes the proof. \square

Lemma 6.14. *Let $M = \langle W, \circ, E \rangle$ be a BBI-model named by ρ and let A be a pure formula. Suppose that $M, w \models_\rho A[\theta]$ for all $w \in W$ and substitutions θ of nominals for nominals. Then A is valid in M .*

Proof. Letting ρ' be a hybrid valuation and $w \in W$, we must show that $M, w \models_{\rho'} A$. Since M is named by ρ , we have that for any $\ell \in \mathcal{N}$ there is a $k \in \mathcal{N}$ such that $\rho(k) = \rho'(\ell)$. Thus we

can define the substitution θ of nominals for nominals by: $\theta(\ell)$ is the first $k \in \mathcal{N}$ with $\rho(k) = \rho'(\ell)$. By hypothesis, we have that $M, w \models_\rho A[\theta]$ for all $w \in W$.

We can now prove by structural induction on A that $M, w \models_{\rho'} A$. In the case that A is a nominal ℓ , we must show that $\rho'(\ell) = w$, and are done since by assumption $w = \rho(\ell[\theta]) = \rho'(\ell)$. Note that A cannot be a propositional variable since it is assumed pure. The other cases follow by induction hypothesis. \square

Theorem 6.15 (Completeness). *Let Ax be a set of pure HyBBI-formulas. Then if a HyBBI-formula is valid in the class of BBI-models satisfying Ax , then it is provable in $\mathbf{K}_{\text{HyBBI}} + Ax$.*

Proof. Suppose A is valid in all BBI-models $M \in \mathcal{C}$, but not provable in $\mathbf{K}_{\text{HyBBI}} + Ax$. Then $\{\neg A\}$ is consistent. Using the Extended Lindenbaum Lemma (6.5), we can construct a named, pasted MCS $\Gamma \supseteq \{\neg A\}$. Now let $M^\Gamma = \langle W^\Gamma, \circ^\Gamma, E^\Gamma \rangle$ be the named model yielded by Γ , and ρ^Γ the corresponding canonical valuation. By Lemma 6.13, M^Γ is a BBI-model.

Furthermore, for any pure formula $B \in Ax$ and any nominal substitution θ , we have that $\vdash B[\theta]$ is provable (using the rule (Subst)), which means that $B[\theta] \in \Delta$ for all $\Delta \in W^\Gamma$ since MCSs are closed under provability. By the Truth Lemma, we obtain

$$M^\Gamma, \Delta \models_{\rho^\Gamma} B[\theta] \quad \text{for all } B \in Ax, \Delta \in W^\Gamma, \text{ and substitutions } \theta$$

Thus, by Lemma 6.14, all formulas in Ax are valid in M^Γ , i.e. $M^\Gamma \in \mathcal{C}$. Thus, by the main assumption, A is valid in M^Γ .

Now, since Γ is named by construction, we have $\Gamma \in W^\Gamma$ by part 4 of Lemma 6.6. Since $\neg A \in \Gamma$, we have $M^\Gamma, \Gamma \models_{\rho^\Gamma} \neg A$ by the Truth Lemma. That is, $M^\Gamma, \Gamma \not\models_{\rho^\Gamma} A$. Thus A is not valid in M^Γ , contradiction. We conclude A is provable in $\mathbf{K}_{\text{HyBBI}} + Ax$ as required. \square

Corollary 6.16. *Let \mathcal{S} be any separation theory from Definition 3.1 not including the cross-split property, and let Ax be the set of pure HyBBI formulas defining \mathcal{S} , as given by Theorem 4.3.*

Then a HyBBI-formula is provable in $\mathbf{K}_{\text{HyBBI}} + Ax$ if and only if it is valid in the class of BBI-models satisfying the separation theory \mathcal{S} .

Proof. Follows from Proposition 5.2, Theorem 6.15 and the fact that the pure formulas Ax collectively define the class of BBI-models satisfying \mathcal{S} , guaranteed by Theorem 4.3. \square

7. HyBBI with the \downarrow binder

In this section, we study the extension of HyBBI with the \downarrow binder from hybrid logic [1–3]. Specifically, we show that the elusive cross-split property from Definition 3.1 is definable in this extension, called HyBBI(\downarrow), and we show how to extend our soundness and parametric completeness results for HyBBI in the previous sections to the setting of HyBBI(\downarrow).

7.1 Formulas and expressivity

Definition 7.1 (HyBBI(\downarrow)-formula). A HyBBI(\downarrow)-formula is defined as a HyBBI-formula (Defn. 4.1), except that if A is a HyBBI(\downarrow)-formula and ℓ a nominal then $\downarrow \ell$. A is also a HyBBI(\downarrow)-formula.

Definition 7.2 (HyBBI(\downarrow)-validity). Given any hybrid valuation ρ for a BBI-model $M = \langle W, \circ, E \rangle$, and any $w \in W$, we extend the definition of the forcing relation for HyBBI in Defn. 4.2 by the following clause for the \downarrow binder:

$$M, w \models_\rho \downarrow \ell. A \Leftrightarrow M, w \models_{\rho[\ell:=w]} A$$

where $\rho[l := w]$ is notation for the hybrid valuation defined as ρ except that $\rho[l := w](\ell) \stackrel{\text{def}}{=} w$.

The definition of validity for HyBBI then extends to HyBBI(\downarrow) in the obvious way.

Proposition 7.3. *The cross-split property (see Definition 3.1) is definable in HyBBI(\downarrow) via the following pure formula:*

$$(a * b) \wedge (c * d) \quad (\text{cs})$$

$$\vdash @_a(\top * \downarrow ac. @_a(\top * \downarrow ad. @_a(ac * ad) \wedge @_b(\top * \downarrow bc. @_b(\top * \downarrow bd. @_b(bc * bd) \wedge @_c(ac * bc) \wedge @_d(ad * bd))))))$$

Proof. We use the following formula abbreviations:

$$\begin{aligned} A &\stackrel{\text{def}}{=} @_a(\top * \downarrow ac. @_a(\top * \downarrow ad. @_a(ac * ad) \wedge B)) \\ B &\stackrel{\text{def}}{=} @_b(\top * \downarrow bc. @_b(\top * \downarrow bd. @_b(bc * bd) \wedge C)) \\ C &\stackrel{\text{def}}{=} @_c(ac * bc) \wedge @_d(ad * bd) \end{aligned}$$

Using the fact that ac , ad and a are distinct nominals, we have for any BBI-model $M = \langle W, \circ, E \rangle$, valuation ρ and $w \in W$,

$$\begin{aligned} &M, w \models_{\rho} @_a(\top * \downarrow ac. @_a(\top * \downarrow ad. @_a(ac * ad) \wedge B)) \\ \Leftrightarrow &M, \rho(a) \models_{\rho} \top * \downarrow ac. @_a(\top * \downarrow ad. @_a(ac * ad) \wedge B) \\ \Leftrightarrow &\rho(a) \in w' \circ w_1 \text{ and} \\ &M, w_1 \models_{\rho} \downarrow ac. @_a(\top * \downarrow ad. @_a(ac * ad) \wedge B) \\ \Leftrightarrow &M, w_1 \models_{\rho[ac:=w_1]} @_a(\top * \downarrow ad. @_a(ac * ad) \wedge B) \\ \Leftrightarrow &M, \rho(a) \models_{\rho[ac:=w_1]} \top * \downarrow ad. @_a(ac * ad) \wedge B \\ \Leftrightarrow &\rho(a) \in w'' \circ w_2 \text{ and} \\ &M, w_2 \models_{\rho[ac:=w_1]} \downarrow ad. @_a(ac * ad) \wedge B \\ \Leftrightarrow &M, w_2 \models_{\rho[ac:=w_1, ad:=w_2]} @_a(ac * ad) \wedge B \\ \Leftrightarrow &M, \rho(a) \models_{\rho[ac:=w_1, ad:=w_2]} (ac * ad) \wedge B \\ \Leftrightarrow &\rho(a) \in w_1 \circ w_2 \text{ and } M, \rho(a) \models_{\rho[ac:=w_1, ad:=w_2]} B \end{aligned}$$

By a similar chain of reasoning, we have

$$\begin{aligned} &M, \rho(a) \models_{\rho[ac:=w_1, ad:=w_2]} B \\ \Leftrightarrow &\rho(b) \in w_3 \circ w_4 \text{ and} \\ &M, \rho(b) \models_{\rho[ac:=w_1, ad:=w_2, bc:=w_3, bd:=w_4]} C \end{aligned}$$

Finally, we have

$$\begin{aligned} &M, \rho(b) \models_{\rho[ac:=w_1, ad:=w_2, bc:=w_3, bd:=w_4]} C \\ \Leftrightarrow &\rho(c) \in w_1 \circ w_3 \text{ and } \rho(d) \in w_2 \circ w_4 \end{aligned}$$

Putting everything together, we have

$$\begin{aligned} &M, w \models_{\rho} A \\ \Leftrightarrow &\exists w_1, w_2, w_3, w_4. \rho(a) \in w_1 \circ w_2, \rho(b) \in w_3 \circ w_4, \\ &\rho(c) \in w_1 \circ w_3, \rho(d) \in w_2 \circ w_4 \end{aligned}$$

With the above equivalence for A in place, we can now show that (cs) defines the cross-split property of Definition 3.1.

(\Rightarrow) Suppose M has the cross-split property. We require to show that the formula (cs) is valid in M , i.e. that if $M, w \models_{\rho} (a * b) \wedge (c * d)$ then $M, w \models_{\rho} A$. Supposing $M, w \models_{\rho} (a * b) \wedge (c * d)$, we have $w \in (\rho(a) \circ \rho(b)) \cap (\rho(c) \circ \rho(d))$. By the cross-split property, there then exist $ac, ad, bc, bd \in W$ such that $\rho(a) \in ac \circ ad$, $\rho(b) \in bc \circ bd$, $\rho(c) \in ac \circ bc$ and $\rho(d) \in ad \circ bd$. Thus, by the equivalence above, $M, w \models_{\rho} A$ as required.

(\Leftarrow) Suppose the formula (cs) is valid in M . We require to show that M has the cross-split property. Suppose $w \in (t \circ u) \cap (v \circ w)$. Define a hybrid valuation ρ for M by

$$\rho(a) = t \quad \rho(b) = u \quad \rho(c) = u \quad \rho(d) = v$$

where a, b, c, d are distinct nominals. We have that $M, w \models_{\rho} (a * b) \wedge (c * d)$. Thus, as (cs) is valid in M , we have $M, w \models_{\rho} A$. Using the equivalence above, there then exist $tv, tw, uv, uw \in W$

such that $t \in tv \circ tw$, $u \in uv \circ uw$, $v \in tv \circ uv$ and $w \in tw \circ uw$ as required. \square

The \downarrow binder of HyBBI(\downarrow) also allows us to encode the definition of the *overlapping conjunction* \wp of separation logic, which has been used in specifying and verifying programs manipulating data structures with intrinsic sharing [15, 25]. In these works, \wp is introduced as a new primitive connective, defined by extending the standard forcing relation for BBI (Definition 2.3) as follows:

$$\begin{aligned} &M, w \models_{\rho} A_1 \wp A_2 \\ \Leftrightarrow &\exists w_1, w_2, w_3, w', w'' \in W. \\ &w' \in w_1 \circ w_2 \text{ and } w'' \in w_2 \circ w_3 \text{ and } w \in w' \circ w_3 \\ &\text{and } M, w' \models_{\rho} A_1 \text{ and } M, w'' \models_{\rho} A_2 \end{aligned}$$

We give below an equivalent formulation of $A_1 \wp A_2$ solely in terms of HyBBI(\downarrow) connectives. We conjecture that this is not possible in BBI (for arbitrary A_1 and A_2).

Proposition 7.4. *For any HyBBI(\downarrow) formulas A_1 and A_2 , the overlapping conjunction $A_1 \wp A_2$ is definable via the following HyBBI(\downarrow) formula, where ℓ and ℓ_s do not appear free in A_1 or A_2 :*

$$\downarrow \ell. \top * \downarrow \ell_s. @_{\ell} \neg(\ell_s \multimap \neg A_1) * \neg(\ell_s \multimap \neg A_2) * \ell_s$$

Proof. Let $M = \langle W, \circ, E \rangle$ be a BBI-model, ρ a valuation for M , and $w \in W$.

$$\begin{aligned} &M, w \models_{\rho} \downarrow \ell. \top * \downarrow \ell_s. @_{\ell} \neg(\ell_s \multimap \neg A_1) * \neg(\ell_s \multimap \neg A_2) * \ell_s \\ \Leftrightarrow &M, w \models_{\rho[\ell:=w]} \top * \downarrow \ell_s. @_{\ell} \neg(\ell_s \multimap \neg A_1) * \neg(\ell_s \multimap \neg A_2) * \ell_s \\ \Leftrightarrow &w \in w' \circ w_s \text{ and} \\ &M, w_s \models_{\rho[\ell:=w]} \downarrow \ell_s. @_{\ell} \neg(\ell_s \multimap \neg A_1) * \neg(\ell_s \multimap \neg A_2) * \ell_s \\ \Leftrightarrow &M, w \models_{\rho[\ell:=w, \ell_s:=w_s]} \neg(\ell_s \multimap \neg A_1) * \neg(\ell_s \multimap \neg A_2) * \ell_s \\ \Leftrightarrow &\exists w_1, w_2, w_3, w'', w'' \in w_1 \circ w_2 \text{ and } w \in w'' \circ w_3 \\ &\text{and } M, w_1 \models_{\rho'} \neg(\ell_s \multimap \neg A_1) \text{ and } M, w_2 \models_{\rho'} \neg(\ell_s \multimap \neg A_2) \\ &\text{and } M, w_3 \models_{\rho'} \ell_s \quad (\text{letting } \rho' = \rho[\ell := w, \ell_s := w_s]) \end{aligned}$$

Notice now that

$$\begin{aligned} &M, w_1 \models_{\rho[\ell:=w, \ell_s:=w_s]} \neg(\ell_s \multimap \neg A_1) \\ \Leftrightarrow &\exists w'_1, w''_1. w'_1 \in w_1 \circ w''_1 \text{ and } M, w'_1 \models_{\rho[\ell:=w, \ell_s:=w_s]} \ell_s \\ &\text{and } M, w'_1 \models_{\rho[\ell:=w, \ell_s:=w_s]} A_1 \\ \Leftrightarrow &\exists w'_1. w'_1 \in w_1 \circ w_s \text{ and } M, w'_1 \models_{\rho[\ell:=w, \ell_s:=w_s]} A_1 \end{aligned}$$

Consequently,

$$\begin{aligned} &M, w \models_{\rho} \downarrow \ell. \top * \downarrow \ell_s. @_{\ell} \neg(\ell_s \multimap \neg A_1) * \neg(\ell_s \multimap \neg A_2) * \ell_s \\ \Leftrightarrow &\exists w_1, w_2, w_s, w'_1, w'_2, w'', w'' \in w_1 \circ w_2 \text{ and } w \in w'' \circ w_s \\ &\text{and } w'_1 \in w_1 \circ w_s \text{ and } w'_2 \in w_2 \circ w_s \\ &\text{and } M, w'_1 \models_{\rho'} A_1 \text{ and } M, w'_2 \models_{\rho'} A_2 \end{aligned}$$

Since \circ is commutative and associative, and ℓ and ℓ_s are sufficiently fresh, this is equivalent to $M, w \models_{\rho} A_1 \wp A_2$. \square

Another variant of the overlapping conjunction was introduced by Cherini and Blanco [10] to deal with some forms of specified sharing. They define an alternative, three-place connective $A_1(\wp : B)A_2$ which tags the shared core of A_1 and A_2 with a formula B that it satisfies. The satisfaction of such a formula is defined as follows:

$$\begin{aligned} &M, w \models_{\rho} A_1(\wp : B)A_2 \\ \Leftrightarrow &\exists w_1, w_2, w_3, w', w'' \in W. \\ &w' \in w_1 \circ w_2 \text{ and } w'' \in w_2 \circ w_3 \text{ and } w \in w' \circ w_3 \\ &\text{and } M, w' \models_{\rho} A_1 \text{ and } M, w'' \models_{\rho} A_2 \text{ and } M, w_2 \models_{\rho} B \end{aligned}$$

It is easy to modify the formula of Proposition 7.4 so as to accommodate this variant. The three-place connective $A_1(\wp : B)A_2$ of [10] is definable using the following HyBBI(\downarrow) formula, parametric in A_1, A_2 and B :

$$\downarrow \ell. \top * \downarrow \ell_s. B \wedge @_{\ell} \neg(\ell_s \multimap \neg A_1) * \neg(\ell_s \multimap \neg A_2) * \ell_s$$

7.2 Proof theory, soundness and completeness

Definition 7.5. We define $\mathbf{K}_{\text{HyBBI}(\downarrow)}$ to be the proof system obtained by adding the following axiom schema to $\mathbf{K}_{\text{HyBBI}}$:

$$(\text{Bind } \downarrow) \quad \vdash @_j(\downarrow \ell. B \leftrightarrow B[j/\ell])$$

Lemma 7.6 (Nominal Substitution Lemma). *We have for any model $M = \langle W, \circ, E \rangle$, hybrid valuation ρ , $\text{HyBBI}(\downarrow)$ -formula A and nominals j, ℓ ,*

$$M, \rho(j) \models_{\rho} A[j/\ell] \Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)]} A$$

where $[j/\ell]$ is a (capture-avoiding) nominal substitution.

Proof. By structural induction on A . The cases not involving nominals are straightforward. We examine the nominal cases, making use of the identity $\rho[\ell := \rho(j)](k) = \rho(k[j/\ell])$:

Case $A = k \in \mathcal{N}$. The required equivalence becomes:

$$\begin{aligned} M, \rho(j) \models_{\rho} k[j/\ell] &\Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)]} k \\ \text{i.e. } \rho(j) = \rho(k[j/\ell]) &\Leftrightarrow \rho(j) = \rho[\ell := \rho(j)](k) \end{aligned}$$

which follows from the identity above.

Case $A = @_k B$. We have $(@_k B)[j/\ell] = @_k [j/\ell] B[j/\ell]$ (noting that $@$ is not a binder, so the nominal substitution applies to the argument k), and proceed as follows:

$$\begin{aligned} M, \rho(j) \models_{\rho} @_k [j/\ell] B[j/\ell] &\Leftrightarrow M, \rho(k[j/\ell]) \models_{\rho} B[j/\ell] \\ &\Leftrightarrow M, \rho[\ell := \rho(j)](k) \models_{\rho} B[j/\ell] \quad (\text{by above identity}) \\ &\Leftrightarrow M, \rho[\ell := \rho(j)](k) \models_{\rho[\ell := \rho(j)]} B \quad (\text{by ind. hyp.}) \\ &\Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)]} @_k B \end{aligned}$$

Case $A = \downarrow k. B$. First we look at the subcase where $k \neq \ell$. In this case, we have $(\downarrow k. B)[j/\ell] = \downarrow k. B[j/\ell]$, and can proceed as follows:

$$\begin{aligned} M, \rho(j) \models_{\rho} \downarrow k. B[j/\ell] &\Leftrightarrow M, \rho(j) \models_{\rho[k := \rho(j)]} B[j/\ell] \\ &\Leftrightarrow M, \rho(j) \models_{\rho[k := \rho(j)][\ell := \rho(j)]} B \quad (\text{by ind. hyp.}) \\ &\Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)][k := \rho(j)]} B \quad (\text{since } k \neq \ell) \\ &\Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)]} \downarrow k. B \end{aligned}$$

In the subcase $k = \ell$, we have $(\downarrow k. B)[j/\ell] = \downarrow \ell. B$ (since ℓ is bound in $\downarrow k. B$), and instead proceed as follows:

$$\begin{aligned} M, \rho(j) \models_{\rho} \downarrow \ell. B &\Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)]} B \\ &\Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)][\ell := \rho(j)]} B \\ &\Leftrightarrow M, \rho(j) \models_{\rho[\ell := \rho(j)]} \downarrow \ell. B \end{aligned}$$

This completes all cases. \square

Proposition 7.7 (Soundness). *Any $\mathbf{K}_{\text{HyBBI}(\downarrow)}$ -provable formula is valid.*

Proof. Given the soundness of $\mathbf{K}_{\text{HyBBI}}$ (Proposition 5.2), we just need to show that the new axiom (Bind \downarrow) is valid in all BBI-models. Let $M = \langle W, \circ, E \rangle$ be a BBI-model, let ρ be a valuation for M and let $w \in W$. We need to show that

$$\begin{aligned} M, w \models_{\rho} @_j((\downarrow \ell. B) \leftrightarrow B[j/\ell]) \\ \text{i.e. } M, \rho(j) \models_{\rho} \downarrow \ell. B \Leftrightarrow M, \rho(j) \models_{\rho} B[j/\ell] \\ \text{i.e. } M, \rho(j) \models_{\rho[\ell := \rho(j)]} B \Leftrightarrow M, \rho(j) \models_{\rho} B[j/\ell] \end{aligned}$$

which is precisely guaranteed by our Nominal Substitution Lemma (that is, Lemma 7.6). \square

We can obtain a parametric completeness result for $\mathbf{K}_{\text{HyBBI}(\downarrow)}$ by repeating the Lindenbaum model construction for $\mathbf{K}_{\text{HyBBI}}$ in section 6. The only point of difference is that the crucial Truth Lemma needs to be extended to account for the \downarrow binder case (cf. [3]).

Lemma 7.8 (Extended Truth Lemma). *For any $\text{HyBBI}(\downarrow)$ -formula A and $\Delta \in W^{\Gamma}$, we have $M^{\Gamma}, \Delta \models_{\rho^{\Gamma}} A$ if and only if $A \in \Delta$.*

Proof. By structural induction on A , with all cases except $A = \downarrow \ell. B$ covered by Lemma 6.12. In this case, using the fact that $\Delta = \{A \mid @_j A \in \Gamma\}$ for some nominal j , we proceed as follows:

$$\begin{aligned} M^{\Gamma}, \Delta \models_{\rho^{\Gamma}} \downarrow \ell. B &\Leftrightarrow M^{\Gamma}, \Delta \models_{\rho^{\Gamma}[\ell := \Delta]} B \\ &\Leftrightarrow M^{\Gamma}, \Delta \models_{\rho^{\Gamma}[\ell := \rho^{\Gamma}(j)]} B \\ &\Leftrightarrow M^{\Gamma}, \Delta \models_{\rho^{\Gamma}} B[j/\ell] \quad (\text{by Lemma 7.6}) \\ &\Leftrightarrow B[j/\ell] \in \Delta \quad (\text{by ind. hyp.}) \\ &\Leftrightarrow @_j B[j/\ell] \in \Gamma \end{aligned}$$

Now since Γ is an MCS and thus closed under $(K_{@})$ and the new axiom (Bind \downarrow), we have $@_j B[j/\ell] \in \Gamma$ if and only if $@_j \downarrow \ell. B \in \Gamma$ if and only if $\downarrow \ell. B \in \Delta$, which completes the case. \square

Theorem 7.9 (Completeness). *Let Ax be any set of pure $\text{HyBBI}(\downarrow)$ -formulas. Then if a $\text{HyBBI}(\downarrow)$ -formula is valid in the class of BBI-models satisfying Ax , then it is provable in $\mathbf{K}_{\text{HyBBI}(\downarrow)} + Ax$.*

Proof. Exactly as Theorem 6.15, using the Extended Truth Lemma (Lemma 7.8) for $\text{HyBBI}(\downarrow)$ in place of Lemma 6.12. \square

Corollary 7.10. *Let \mathcal{S} be any separation theory from Definition 3.1, and let Ax be the set of pure $\text{HyBBI}(\downarrow)$ formulas defining the properties \mathcal{S} , as given by Theorem 4.3 and Proposition 7.3.*

Then a $\text{HyBBI}(\downarrow)$ -formula is provable in $\mathbf{K}_{\text{HyBBI}(\downarrow)} + Ax$ if and only if it is valid in the class of BBI-models satisfying the separation theory \mathcal{S} .

Proof. Follows from Proposition 7.7, Theorem 7.9 and the definability results in Theorem 4.3 and Proposition 7.3. \square

8. Conclusions and future work

In this paper, we show that many separation theories that arise naturally in applications of separation logic are not definable in the standard propositional basis for separation logic, namely BBI. To overcome these limitations in expressivity, we introduce new hybrid versions of BBI, obtained by marrying BBI with the machinery of hybrid logics. In addition, we show how to obtain axiomatic proof systems for these hybrid logics that are sound and complete for any separation theory obtained by combining properties from a list of those we found in the separation logic literature.

In future work, we plan to explore possible applications of our hybrid logics to program analysis, e.g. by adding support for nominals to separation logic. This would allow for more expressive program specifications, and for instance, as a conjecture, would capture the immutable specs introduced by David and Chin [11]. In these specifications, some parts of the heap are tagged as immutable using the syntax $A@I$, as in the following specification of a function that computes the length of a linked list given as argument:

$$\{list(x, n)@I\} \text{length}(x) \{ret = n\}$$

where $list(x, n)$ is an (inductive) predicate describing a linked list of length n starting at address x , and ret is the return value of the function. Notice that immutable parts of the heap are not repeated

in the post-condition; their presence is implicit. We could represent such a triple using nominals as

$$\{list(x, n) \wedge \ell\} \text{length}(x) \{\ell \wedge ret = n\}$$

which stipulates that the heap ℓ of the precondition is found intact in the post.

More broadly, we hope that our introduction of more expressive intermediaries between BBI and full first-order logic will help facilitate the expression and verification of more complex program properties, particularly those involving overlapping data structures.

Acknowledgments

Many thanks to Hongseok Yang for pointing out to us the need for Lemma 6.10.

References

- [1] Patrick Blackburn, Maarten de Rijke, and Yde Venema. *Modal Logic*. Cambridge University Press, 2001.
- [2] Patrick Blackburn and Jerry Seligman. Hybrid languages. *Journal of Logic, Language and Information*, 4(3):251–272, 1995.
- [3] Patrick Blackburn and Balder ten Cate. Pure extensions, proof rules, and hybrid axiomatics. *Studia Logica*, 84(2):277–322, 2006.
- [4] George E.P. Box and Norman R. Draper. *Empirical model-building and response surfaces*. Wiley series in probability and mathematical statistics. John Wiley & Sons, Inc., 1987.
- [5] James Brotherston and Cristiano Calcagno. Classical BI: Its semantics and proof theory. *Logical Methods in Computer Science*, 6(3), 2010.
- [6] James Brotherston and Max Kanovich. Undecidability of propositional separation logic and its neighbours. Submitted, 2012.
- [7] Cristiano Calcagno, Dino Distefano, Peter O’Hearn, and Hongseok Yang. Compositional shape analysis by means of bi-abduction. *Journal of the ACM*, 58(6), December 2011.
- [8] Cristiano Calcagno, Philippa Gardner, and Uri Zarfaty. Context logic as modal logic: Completeness and parametric inexpressivity. In *Proceedings of POPL-34*, pages 123–134. ACM, 2007.
- [9] Cristiano Calcagno, Peter O’Hearn, and Hongseok Yang. Local action and abstract separation logic. In *Proceedings of LICS-22*, pages 366–378. IEEE Computer Society, 2007.
- [10] Renato Cherini and Javier O. Blanco. Local reasoning for abstraction and sharing. In Sung Y. Shin and Sascha Ossowski, editors, *SAC*, pages 552–557. ACM, 2009.
- [11] Cristina David and Wei-Ngan Chin. Immutable specifications for more concise and precise verification. In *Proceedings of OOPSLA-11*, pages 359–374. ACM, 2011.
- [12] Thomas Dinsdale-Young, Lars Birkedal, Philippa Gardner, Matthew J. Parkinson, and Hongseok Yang. Views: compositional reasoning for concurrent programs. In Roberto Giacobazzi and Radhia Cousot, editors, *POPL*, pages 287–300. ACM, 2013.
- [13] Robert Dockins, Aquinas Hobor, and Andrew W. Appel. A fresh look at separation algebras and share accounting. In *Proceedings of APLAS*, volume 5904 of *LNCS*, pages 161–177. Springer-Verlag, 2009.
- [14] Didier Galmiche and Dominique Larchey-Wendling. Expressivity properties of Boolean BI through relational models. In *Proceedings of FSTTCS*, volume 4337 of *LNCS*, pages 357–368. Springer, 2006.
- [15] Aquinas Hobor and Jules Villard. The ramifications of sharing in data structures. In *POPL*, pages 523–536. ACM, 2013.
- [16] Zhe Hou, Alwen Tiu, and Rajeev Goré. A labelled sequent calculus for Boolean BI: Proof theory and proof search. Submitted, 2013.
- [17] Samin Ishtiaq and Peter W. O’Hearn. BI as an assertion language for mutable data structures. In *Proceedings of POPL-28*, pages 14–26. ACM, 2001.
- [18] Richard Kaye. *Models of Peano Arithmetic*. Oxford University Press, 1991.
- [19] Dominique Larchey-Wendling. The formal strong completeness of Boolean BI. Submitted, 2012.
- [20] Dominique Larchey-Wendling and Didier Galmiche. Exploring the relation between intuitionistic BI and Boolean BI: An unexpected embedding. *Mathematical Structures in Computer Science*, 19:1–66, 2009.
- [21] Dominique Larchey-Wendling and Didier Galmiche. The undecidability of Boolean BI through phase semantics. In *Proceedings of LICS-25*, pages 140–149. IEEE Computer Society, 2010.
- [22] Jonghyun Park, Jeongbong Seo, and Sungwoo Park. A theorem prover for Boolean BI. In *Proceedings of POPL-40*, pages 219–232. ACM, 2013.
- [23] David Pym. *The Semantics and Proof Theory of the Logic of Bunched Implications*. Applied Logic Series. Kluwer, 2002. Errata and remarks (Pym 2008) maintained at <http://www.abdn.ac.uk/~csc335/BI-monograph-errata.pdf>.
- [24] John C. Reynolds. Separation logic: A logic for shared mutable data structures. In *Proceedings of LICS-17*, pages 55–74. IEEE Computer Society, 2002.
- [25] John C. Reynolds. A short course on separation logic. 2003.
- [26] H. Yang, O. Lee, J. Berdine, C. Calcagno, B. Cook, D. Distefano, and P. O’Hearn. Scalable shape analysis for systems code. In *Proceedings of CAV-20*, volume 5123 of *LNCS*, pages 385–398. Springer, 2008.